



The miracles of science™

SPEED, AGILITY AND EXCEPTIONAL COVERAGE

DuPont™
Acapela®
fungicide

Protect your soybean and corn crops against key diseases.

DuPont™ Acapela® is a broad-spectrum fungicide designed to provide reliable protection against Northern corn leaf blight in corn and sclerotinia rot (white mould), Asian soybean rust, frogeye leafspot and septoria brown spot in soybeans.

With 4 movement properties, Acapela® is a one-of-a-kind fungicide that quickly and efficiently surrounds, penetrates and protects the leaf and stem.

This characteristic provides superior disease control for greener, healthier crops and higher yield potential.



Translaminar Movement

Moves through the leaf surface to protect top and bottom sides of the leaf.



Local Gas Activity

Gas-like protective barrier that moves locally over the leaf surface.



Xylem Systemic Activity

Movement through the plant tissues to distribute throughout the leaf.



Wax Diffusion


Provides more consistent coverage across the leaf or stem surface.




Rapid Absorption: Acapela® is also rapidly absorbed, putting these 4 unique movement properties to work fast, and allows you to spray when field conditions are challenging.



Translaminar Movement



Xylem Systemic Activity




Local Gas Activity



Wax Diffusion Activity


More moves, more bushels

Acapela® Field Performance - Soybean



Source: 2013 - 2015 DuPont Side-by-Side Trials

Acapela® Field Performance - Corn VT-R3
(Standard Rate 40 acres/jug)



Source: 2013 DuPont side-by-side trials (ON and QC)

Crops	Diseases	Rates	Application Information
Corn	Northern corn leaf blight	0.213 to 0.325 L/ac (30 to 45 acres/jug)	Begin applications prior to disease development and continue on a 7 to 14 day interval. Use higher rate and shorter interval when disease pressure is high. For optimal disease control, apply at full tassel (VT) to milk stage (R3) corn.
Soybeans	Sclerotinia rot – white mould ¹	0.352 L/ac (27 acres/jug)	For white mould: make initial preventive application at beginning bloom and follow with 2nd application with a different mode of action 7 to 10 days later at full bloom.
	Asian soybean rust Frogeye leafspot Septoria brown spot	0.240 to 0.352 L/ac (27 to 40 acres/jug)	Apply at R2 (full bloom) stage of development.

Acapela® Helps Optimize Soybean Yields



Source: Rosser, MB, 2015

Other application information

Packaging	One (1) case = 2 x 9.6 L jugs Acapela® is also available in a 115.2 L drum
Water Volumes	Ground: 110 L/ha = 10 gal/ac. Air: 50 L/ha = 4.5 gal/ac.
Aerial Application	Acapela® is registered for aerial application.

For more information on Acapela®, contact your local DuPont representative, call the DuPont™ FarmCare® Support Centre at 1-800-667-3925 or visit acapela.dupont.ca

¹Suppression

As with all crop protection products, read and follow label instructions carefully.

All information effective December 2015.
Member of CropLife Canada.

Unless indicated, trademarks with ®, ™ or SM are trademarks of DuPont or affiliates. © 2016 DuPont.
Quilt®, Allegro® (Syngenta), Delaro™ (Bayer CropScience), Priaxor® (BASF)



The miracles of science™