



CLEANER FIELDS, HIGHER YIELDS

DuPont™
Sortan™ IS
herbicide
Pre-emergence or post-emergence

New Sortan™ IS will allow growers to remove early season weed pressure, including volunteer Roundup Ready® canola, which is key to maximizing yield and profits in corn.

Removes
Early-Season Weed
Competition

Designed to remove **early season weed competition** including volunteer Roundup Ready® canola and wild buckwheat

Resistance
Management

Provides an **additional mode of action** to glyphosate, making it a great tool for effective resistance management

Extended Control

Extended control of tough broadleaf and grassy weeds throughout the critical weed free period

Application Timing

Sortan™ IS can be used as a pre-emergent or post-emergent application (spike to V3 stage). Recommended to tank-mix with a glyphosate herbicide at 365 g ai/acre (900 ai g/ha) for control of additional weeds (glyphosate-tolerant corn only).

Research trials show Sortan™ IS provides outstanding control of volunteer canola*



Sortan™ IS

Glyphosate alone

Extended control of Wild Buckwheat



Armezon® + Aatrex® Liquid 480+
glyphosate post-emerge

Sortan™ IS + glyphosate
post-emerge

*Tank-mix with glyphosate required for volunteer Clearfield® canola control.
Source: Carman, 2015

Morris, MB: August 9, 2016, 63 DAA

Crop Rotation


The following crops may be seeded 10 months after application: Spring wheat (including durum), oats, barley, canola, soybeans, dry beans, chickpeas, potatoes, sunflowers, field corn, and field peas.

Application Information		Weeds Controlled
Pre-Emergent Rate	30 g/acre, 40 acres/bottle	Volunteer Canola*
Post-Emergent Rate	15 g/acre – 30 g/acre, 40 – 80 acres/bottle (Refer to product label for more details on the post-emergent application).	Wild Buckwheat
Packaging	1.2 kg bottle	Redroot Pigweed (including triazine resistant)
Tank-Mix	Tank-mix with a glyphosate herbicide at 365 g ai/acre (900 g ai/ha) for control of additional weeds (glyphosate-tolerant corn only).	Lamb's Quarters
Water Volume	Minimum 40 L/acre (9 imp. gal/acre)	Barnyard Grass
		Green Foxtail
		Yellow Foxtail ¹
		Quackgrass
		Large (Hairy) Crabgrass ¹
		Fall Panicum
		Old Witchgrass
		¹ Suppression only.
		*Tank-mix with glyphosate required for volunteer Clearfield® canola control.
		Refer to product label for complete details prior to use.

Application Timing

Pre-Emergent application - before emergence of crop or weeds

Post-Emergent application - VE (spike) to V3 (5 leaf stage[†])



[†]Corn leaf stages can vary depending on both growing conditions and corn seed product.

Critical Weed Free Period

New Sortan™ IS is a powerful and versatile tool that can be applied from pre-emergence or spike to the V3 stage, which will help growers keep their corn weed free during the critical weed-free period.

- Critical weed free period in corn is from emergence to the V4 (6 leaf) stage.
- Removing weeds during this period is key to minimizing yield loss due to weed competition in corn.

As with all crop protection products, read and follow label instructions carefully.
Member of CropLife Canada.

Copyright © 2017 DuPont. All rights reserved. The DuPont Oval Logo, DuPont™, and all products denoted with ® or ™ are trademarks or registered trademarks of E. I. du Pont de Nemours and Company or its affiliates. Armezon® (BASF), RoundUp Ready® (Monsanto), Aatrex® Liquid 480 (Syngenta)

