

PMRA APPROVED LABEL, JB
2014-09-16, Sub. No. 2014-3535

[[PRIMARY DISPLAY PANEL]]

GROUP	2, 14	HERBICIDE
--------------	--------------	------------------

Guardian® Plus WDG Herbicide

Wettable granules

FOR SALE FOR USE ON SOYBEANS IN EASTERN CANADA ONLY

COMMERCIAL

REGISTRATION NO. 31494 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND THIS BOOKLET BEFORE USING

GUARANTEE: Chlorimuron-ethyl 5.14%
Flumioxazin 40.59%

Warning: This product contains the allergen sulfite.

NET CONTENTS: 176g -bulk

E. I. DU PONT CANADA COMPANY
AGRICULTURAL PRODUCTS
P.O. BOX 2300, STREETSVILLE
MISSISSAUGA, ONTARIO
L5M 2J4
1-800-667-3925

Skull&crossbones-triangle
CAUTION - POISON

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN.
Avoid breathing spray mist.
Avoid contact with skin, eyes and clothing.
Keep the outer container or bag and soluble bags away from water.
Do not contaminate any body of water.
Harmful if inhaled.

FIRST AID:

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED: Move the person to fresh air. If the person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

For medical emergencies call 1-800-441-3637 (24 hours).

TOXICOLOGICAL INFORMATION: Treat symptomatically

STORAGE:

Store product in original container in a secure, dry area away from food or feed. Not for use or storage in or around the home. Keep container tightly closed.

DISPOSAL:

For recyclable containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use. If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For returnable containers:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on this label. It is an offence under the *PEST CONTROL PRODUCTS ACT* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

[[DIRECTIONS FOR USE LABEL]]

GROUP

2, 14

HERBICIDE

Guardian® Plus WDG Herbicide

Wettable granules

FOR SALE FOR USE ON SOYBEANS IN EASTERN CANADA ONLY

COMMERCIAL

REGISTRATION NO. 31494 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND THIS BOOKLET BEFORE USING

GUARANTEE: Chlorimuron-ethyl 5.14%
Flumioxazin 40.59%

Warning: This product contains the allergen sulfite.

NET CONTENTS: 176g - bulk

E. I. DU PONT CANADA COMPANY
AGRICULTURAL PRODUCTS
P.O. BOX 2300, STREETSVILLE
MISSISSAUGA, ONTARIO
L5M 2J4
1-800-667-3925

Skull&crossbones-triangle

CAUTION - POISON

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN.
Avoid breathing spray mist.
Avoid contact with skin, eyes and clothing.
Keep the outer container or bag and soluble bags away from water.
Do not contaminate any body of water.
Harmful if inhaled.

FIRST AID:

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED: Move the person to fresh air. If the person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

For medical emergencies call 1-800-441-3637 (24 hours).

IMPORTANT:

Injury to or loss of desirable trees or vegetation may result from failure to observe the following:

Do not apply, drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, tennis courts or similar areas. Prevent drift of spray to desirable plants. Do not contaminate any body of water, including irrigation water that may be used on other crops. Carefully observe sprayer clean-up instructions, as spray tank residue may damage crops other than soybeans.

DO NOT APPLY BY AIR.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.
Do not eat, drink or smoke during work. Wash hands and face thoroughly before eating, drinking, smoking, chewing gum, or using the toilet.

Immediately wash off accidental splashes of the concentrate or spray mixture from skin, clothing and out of eyes. Remove clothing immediately if pesticide gets inside. Wash thoroughly and put on clean clothing. After work, change clothing and wash entire body thoroughly. Wash contaminated working clothes separately from other laundry before reuse.

Follow mixer/loader and applicator scenario, as appropriate in the chart below. In addition, wear coveralls over long-sleeved shirt and long pants, chemical-resistant gloves, socks and shoes, goggles or faceshield, during clean-up and repair activities.

Equipment	Personal Protective Equipment		Maximum amount of product handled per day (kg)
	Mixer/Loader	Applicator	
Groundboom	Chemical-resistant coveralls and chemical-resistant gloves, socks and shoes, and goggles or faceshield	Open cab, coveralls over long-sleeved shirt, long pants, socks and shoes, and chemical-resistant gloves, and respirator with a NIOSH/MSHA/BHSE approved organic vapour-removing cartridge with a prefilter approved for pesticides, or a NIOSH/MSHA/BHSE approved canister approved for pesticides	55 kg
		Closed cab, long-sleeved shirt, long pants, socks and shoes. (No gloves are required, but must be present in the cab for maintenance)	

Do not apply when weather conditions favour spray drift from treated areas. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Keep all unprotected persons out of operating areas, or vicinity where there may be drift. Only protected handlers may be in the area during application.

Do not enter or allow worker entry into treated areas until the restricted entry interval of 12 hours for all crop uses.

Do not apply to fine-textured soils.

Do not graze treated fields or feed treated forage or hay to livestock.

Do not apply within 100 metres of non-dormant pears.

Do not apply to powdery soils or soils that are susceptible to wind displacement unless irrigation can be applied immediately after application.

TOXICOLOGICAL INFORMATION: Treat symptomatically

ENVIRONMENTAL HAZARDS :

This product is toxic to aquatic organisms and non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE. Toxic to small wild mammals. Toxic to certain beneficial insects. Minimize spray drift to reduce harmful effects on beneficial insects in habitats next to the application site such as hedgerows and woodland.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

GENERAL INFORMATION:

GUARDIAN® PLUS WDG Herbicide is a pre-plant or post-plant pre-emergence soybean herbicide that will control or suppress select grassy and broadleaf weeds. GUARDIAN® PLUS WDG Herbicide must be applied pre-emergent to the main flush of actively growing weeds. When tank-mixed with a glyphosate herbicide as part of a pre-plant or post-plant burndown, emerged weeds will also be controlled. GUARDIAN® PLUS WDG Herbicide may only be applied using ground equipment. For best results, ensure thorough spray coverage of target weeds if present at the time of application.

Degree of control and duration of effect depend on weed sensitivity, weed size, growing conditions and spray coverage.

DIRECTIONS FOR USE:

GUARDIAN® PLUS WDG Herbicide offers residual control of susceptible broadleaf and grass weeds listed on this label and assists in the control of acetolactate synthase (ALS) resistant weeds. The length of residual control is dependent on rainfall and temperature conditions. Length of residual control will decrease as temperature and precipitation increase and on soils of high organic matter and/or high clay content.

Crop injury may occur from applications made to poorly drained soil and/or applications made under cool, wet conditions. Severe crop injury will result when soils are flooded following applications of GUARDIAN® PLUS WDG Herbicide. Risk of crop injury can be minimized by using on well drained soils, planting at least 4 cm deep, using high quality seed and completely covering seeds with soil prior to pre-emergence applications. Treated soil that is splashed onto newly emerged crops may result in temporary crop injury.

Moisture is necessary to activate GUARDIAN® PLUS WDG Herbicide in soil for residual weed control. Dry weather following applications of GUARDIAN® PLUS WDG Herbicide may reduce effectiveness. However, when adequate moisture is received after dry conditions, GUARDIAN® PLUS WDG Herbicide will control susceptible germinating weeds. GUARDIAN® PLUS WDG Herbicide may not control weeds that germinate after application but before an activating rainfall/irrigation or weeds that germinate through cracks resulting from dry soil.

No-till planters that incorporate the soil during planting may result in decreased weed control in the row. When these types of planters are used, apply GUARDIAN® PLUS WDG Herbicide within 3 days after planting and before soybeans emerge.

Do not perform any tillage operations after application or weed control will be reduced. Apply only once during a single growing season.

RATES AND TIMING:

GUARDIAN® PLUS WDG Herbicide may be applied to soybeans prior to planting or within 3 days after planting and prior to soybean emergence. Application after the soybeans have begun to crack, or are emerged, will result in severe crop injury.

GUARDIAN® PLUS WDG Herbicide provides pre-emergence control of susceptible weeds in soybeans. Apply GUARDIAN® PLUS WDG Herbicide with ground equipment before planting, during planting, or after planting, but before the crop emerges. Do not apply by air. Apply using ground application equipment only.

GUARDIAN® PLUS WDG HERBICIDE APPLIED ALONE

GUARDIAN® PLUS WDG Herbicide can be applied pre-plant or post-plant pre-emergence to soybeans. GUARDIAN® PLUS WDG Herbicide must be applied pre-emergent to the main flush of actively growing weeds.

Apply 176 g/ha of GUARDIAN® PLUS WDG Herbicide prior to the emergence of weeds. Do not apply to fine-textured soils.

WEEDS CONTROLLED BY GUARDIAN® PLUS WDG HERBICIDE ALONE

WEEDS
Canada fleabane
Common chickweed
Common ragweed
Dandelion *
Eastern black nightshade
Fall panicum **
Green foxtail **
Green pigweed
Hairy nightshade
Lamb's-quarters
Redroot pigweed

* Early-season control only on medium-textured soils.

**Suppression only. Weed suppression is a visual reduction in weed competition (reduced population or vigour) as compared to an untreated area. Degree of suppression will vary with environmental conditions prior to and following treatment.

GUARDIAN® PLUS WDG HERBICIDE PLUS GLYPHOSATE:

If weeds are present at the time of application, tank mix GUARDIAN® PLUS WDG Herbicide with a glyphosate herbicide for the burndown of emerged weeds. Apply the tank mix pre-plant or post-plant pre-emergence to soybeans, in conventional or glyphosate tolerant varieties.

Apply 176 g/ha of GUARDIAN® PLUS WDG Herbicide, plus a glyphosate herbicide (present as the potassium, isopropylamine, or di-ammonium salt) at 900 g ae/ha, (e.g., 2.5 L/ha for herbicides containing 360 g ae/L glyphosate or 1.67 L/ha for herbicides containing 540 g ae/L glyphosate).

WEEDS CONTROLLED BY GUARDIAN® PLUS WDG HERBICIDE PLUS GLYPHOSATE

GUARDIAN® PLUS WDG Herbicide plus a glyphosate herbicide will control emerged weeds, including annual sowthistle, velvetleaf, and yellow nutsedge, as well as provide residual control or suppression of all weeds listed for GUARDIAN® PLUS WDG Herbicide applied alone. Consult the respective glyphosate herbicide label for specific instructions for 'Directions for Use', Precautions, and for other weeds controlled by glyphosate herbicide.

When applied as a tank-mix combination, read and observe all label directions, including rates, restrictions and grazing limitations for each product used in the tank-mix. Follow the more stringent label precautions measures for mixing, loading and applying stated on both product labels.

MIXING INSTRUCTIONS:

1. Fill clean tank about 1/4 full with fresh water.
2. Turn on full agitation.
3. Add the proper amount of GUARDIAN® PLUS WDG Herbicide to the water in the spray tank with the agitator running. Continuous agitation is required to keep the product in suspension.
4. After GUARDIAN® PLUS WDG Herbicide has been well mixed and is in suspension, add the glyphosate tank mix partner.
5. Fill the remainder of the spray tank with clean water.

On repeat tank loads, ensure that the amount of spray solution left in the tank from the previous load is less than 10% of the volume about to be mixed.

Agitation is required for uniform mixing and application. The optimum water volume for GUARDIAN® PLUS WDG Herbicide application is 140-190 litres of water per hectare (minimum of 100 litres of water per hectare). For the tank-mix of GUARDIAN® PLUS WDG Herbicide with glyphosate, the recommended water volume is 100-120 litres per hectare. Flat fan nozzles are recommended. Use 50 mesh filter screens or larger. Use spray preparation of GUARDIAN® PLUS WDG Herbicide within 24 hours or product degradation may occur resulting in a loss of weed control. Use vigorous agitation to thoroughly disperse spray mixtures that have been allowed to stand in the tank.

NOTE: GUARDIAN® PLUS WDG Herbicide will degrade in acidic or highly alkaline water. Mix no more than can be used in one day. If spraying is interrupted, thoroughly re-agitate the spray mixture before resuming spraying.

Do not mix or load where there is potential to contaminate wells or aquatic systems.

Field sprayer application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. Boom height must be 60 cm or less above the crop or ground.

DO NOT apply by air.

Use caution when applying under circumstances where possible drift to unprotected persons or food, forage or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption can occur.

Use the largest droplet size consistent with acceptable efficacy. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much pressure. Make application when the wind velocity favours on-target product deposition.

Do not make applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with increasing distance above the ground. Mist or fog may indicate the presence of an inversion in humid areas. Where permissible by local regulations, the applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

BUFFER ZONES:

Overspray or drift to important aquatic and terrestrial wildlife habitats should be avoided. Leave a 16 metre buffer zone between the last spray swath and bodies of water such as ponds, lakes, rivers, and streams. Leave a 39 metre buffer zone between the last spray swath and wetlands, vegetated areas at the edges of bodies of water, woodlots and shelterbelts, and other terrestrial wildlife habitats.

SPRAYER CLEANUP:

To avoid subsequent injury to crops other than soybeans immediately after spraying and prior to spraying other crops, thoroughly remove all traces of GUARDIAN® PLUS WDG Herbicide from mixing and spray equipment as follows:

1. Drain tank; then flush tank, boom and hoses with clean water for a minimum of ten minutes. Visually inspect tank to assure removal of all visible residues of GUARDIAN® PLUS WDG Herbicide. If necessary, repeat step 1. DO NOT CLEAN SPRAY EQUIPMENT NEAR WELLS, AQUATIC SYSTEMS, OR OTHER WATER SOURCES, OR NEAR DESIRABLE VEGETATION.
2. Fill tank with clean water while adding 1 litre household ammonia (containing a minimum 3% ammonia) per 100 litres of water. Flush solution through boom and hoses, and then add more water to completely fill tank. Allow to sit for 15 minutes with agitation. Again flush the hoses, boom and nozzles with the cleaning solution and drain tank.

3. Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.
4. Repeat Step 2.
5. Thoroughly rinse the tank with clean water for a minimum of 5 minutes, flushing water through the hoses and boom.

REPLANTING TO OTHER CROPS:

GUARDIAN® PLUS WDG Herbicide is degraded by natural soil processes, and field tests have shown that the following crops may safely be planted at the prescribed interval following soybeans that have been treated with GUARDIAN® PLUS WDG Herbicide. The recropping intervals are dependent on the pH of the soil, as well as the sensitivity of the specific rotational crop.

Replanting Interval (Months)

Soil pH	Winter Wheat	Field Corn, Soybeans	Alfalfa
≤7.0	4	10	11
>7.0≤7.4	4	10	11
>7.4≤7.8	4	10	Do Not Plant
>7.8≤8.0	4	Do Not Plant	

For other crops, a field bioassay is recommended before planting. A successful field bioassay means growing to maturity a test strip of the crop(s) intended for production the following year.

FIELD BIOASSAY:

Select a representative area or areas of the field previously treated with GUARDIAN® PLUS WDG Herbicide to plant your bioassay crop(s). Be sure to consider factors such as size of field, soil texture, drainage and turn-around areas when selecting the site(s) that are most representative of the soil conditions in the field. On large fields, more than one site may be needed in order to obtain reliable results.

Plant the test strips perpendicular to the direction in which the field was sprayed. The strips should be long enough to cross the width of several spray swaths. Large test strip areas are more reliable than small ones.

Use standard tillage and seeding equipment to plant the bioassay. Prepare a seed bed and plant the crops and varieties you want the option of growing the following year. It is important to use the same planting time, conditions, techniques and cultural practices you normally use to plant and grow the bioassay crop(s). Also plant into an adjacent area not treated with GUARDIAN® PLUS WDG Herbicide to use as a comparison.

As the crop(s) emerges and grows, examine these key points in GUARDIAN® PLUS WDG Herbicide treated and non-treated areas:

- crop stand
- plant colour and vigour
- root development
- yield
- rate of growth

Allow the bioassay crop(s) to grow to maturity while making your observations. Do not overspray the test strips with herbicides that may damage the bioassay crop(s). If the bioassay indicates that GUARDIAN® PLUS WDG Herbicide residues are still present, continue cropping only to those crops listed on the label and do not rotate to other crops until bioassay results indicate that susceptible crops are growing normally.

USE PRECAUTIONS:

Do not apply GUARDIAN® PLUS WDG Herbicide during periods of intense rainfall or to soils saturated with water. Do not apply directly to standing or running water. Do not apply in areas where surface water from the treatment site can run off to adjacent cropland, either planted or to be planted, or into bodies of water (such as streams, ponds, rivers, lakes, and irrigation water), wetlands, or wells. Applications should only be made when there is no hazard of spray drift contaminating non-target land areas since very small quantities of the GUARDIAN® PLUS WDG Herbicide spray solution may severely injure susceptible crops and other sensitive terrestrial habitats during both growing and dormant periods.

Under extreme weather conditions, such as hot, dry weather, excessive moisture, or frost, weed control may be reduced.

Rainfall within 4 hours after application of GUARDIAN® PLUS WDG Herbicide may reduce weed control.

Overspray or drift to important aquatic and terrestrial wildlife habitats should be avoided. Leave a 16 metre buffer zone between the last spray swath and bodies of water such as ponds, lakes, rivers, and streams. Leave a 39 metre buffer zone between the last spray swath and wetlands, vegetated areas at the edges of bodies of water, woodlots and shelterbelts, and other terrestrial wildlife habitats.

GUARDIAN® PLUS WDG Herbicide must not be applied within 60 days of harvest.

Make only one application of GUARDIAN® PLUS WDG Herbicide per season.

Do not graze the treated crop or cut for hay; sufficient data is not available to support such use.

Do not apply GUARDIAN® PLUS WDG Herbicide to soybean grown on soil of pH \geq 8.0 due to risk of rotational crop injury.

RESISTANCE-MANAGEMENT RECOMMENDATIONS

For resistance management, please note that GUARDIAN® PLUS WDG Herbicide is both a Group 2 and Group 14 herbicide. Any weed population may contain or develop plants naturally resistant to Group 2 and/or Group 14 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- ◆ Where possible, rotate the use of GUARDIAN® PLUS WDG Herbicide or other Group 2 and/or Group 14 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- ◆ Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- ◆ Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- ◆ Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- ◆ Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.

- ◆ Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- ◆ For further information or to report suspected resistance, contact your local E. I. du Pont Canada Company representative or the E. I. du Pont Canada Company hotline at 1-800-667-3925.

STORAGE AND DISPOSAL:

STORAGE:

Store product in original container in a secure, dry area away from food or feed. Not for use or storage in or around the home. Keep container tightly closed.

DISPOSAL:

For recyclable containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use. If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For returnable containers:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on this label. It is an offence under the *PEST CONTROL PRODUCTS ACT* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

®/™ Registered trademarks/trademark of E. I. du Pont Company. E. I. du Pont Canada Company is a licensee. Member of CropLife.

* All other products mentioned are trademarks of their respective companies.