

MUSTER® Toss-N-Go® HERBICIDE

75% DRY FLOWABLE

FOR SALE FOR USE ON CANOLA (RAPESEED) AND CONDIMENT MUSTARD (BROWN AND ORIENTAL MUSTARDS ONLY) IN THE PRAIRIE PROVINCES AND PEACE RIVER REGION OF BRITISH COLUMBIA; FOR SALE FOR USE IN CANOLA IN THE OKANAGAN AND CRESTON VALLEY AREAS OF BRITISH COLUMBIA, SOUTHERN ONTARIO, QUEBEC AND THE MARITIMES ONLY; FOR SALE FOR USE ON ORIENTAL MUSTARD (CONDIMENT AND OIL SEED TYPES) IN WESTERN CANADA ONLY. ALSO FOR SALE FOR USE ON SUNFLOWER, LAURENTIAN RUTABAGA AND BRASSICA CARINATA

AGRICULTURAL

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

GUARANTEE: Ethametsulfuron-methyl 75%

Warning, contains the allergen sulfites.

REGISTRATION NO.: 23569 PEST CONTROL PRODUCTS ACT

DO NOT SELL WATER SOLUBLE BAGS SEPARATELY.

NET CONTENTS: 80-800 GRAMS
(1-10 x 80 gram water soluble bags)

E. I. DU PONT CANADA COMPANY
AGRICULTURAL PRODUCTS
P.O BOX 2300, STREETSVILLE
MISSISSAUGA, ONTARIO
L5M 2J4
1-800-667-3925

PRECAUTIONS:

- **KEEP OUT OF REACH OF CHILDREN.**
- Avoid breathing spray mist.
- Avoid contact with skin, eyes and clothing.
- Do not contaminate any body of water.
- Wear chemical resistant gloves during mixing, loading and cleanup.
- Wash hands and face before eating, drinking, smoking and using the toilet.
- Take a shower immediately after use.
- Wash protective clothing in detergent and hot water, separately from household laundry, before reuse.
- **DO NOT APPLY BY AIR.**
- When applied as a tank-mix combination, read and observe all label directions, including rates, restrictions, and grazing limitations for each product used in the tank-mix. Follow the more stringent label precautionary and PPE measures for mixing, loading and applying stated on both product labels.
- Wear long-sleeved shirt, long pants, socks and chemical-resistant gloves and footwear during mixing, loading, application, clean-up and repair. Chemical-resistant gloves are not required while operating groundboom sprayers.

ENVIRONMENTAL HAZARDS

TOXIC to aquatic organisms and non target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE.

Runoff:

To reduce runoff from treated areas into aquatic habitats, consider the characteristics and conditions of the site before treatment. Site characteristics and conditions that may lead to runoff include, but are not limited to: heavy rainfall, moderate to steep slope, bare soil, poorly draining soil (e.g. soils that are compacted or fine textured such as clay).

Avoid application of this product when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

Leaching:

The use of this chemical may result in contamination of groundwater, particularly in areas where soil pH>7, are permeable (e.g. sandy soil) and/or the depth to the water table is shallow.

FIRST AID:

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance,

then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

For medical emergencies call 1-800-441-3637 (24 hours).

TOXICOLOGICAL INFORMATION: Treat symptomatically.

STORAGE:

Store product in original container only, away from fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed. Keep water soluble bags in original outer bag away from moisture.

DISPOSAL:

1. Make the empty container unsuitable for further use.
2. Dispose of the container in accordance with provincial requirements.
3. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the PEST CONTROL PRODUCTS ACT to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

The seller warrants that the purchase by the buyer and the use of this product, as such, will not infringe any Canadian patent.

MUSTER® Toss-N-Go® HERBICIDE

75% DRY FLOWABLE

FOR SALE FOR USE ON CANOLA (RAPESEED) AND CONDIMENT MUSTARD (BROWN AND ORIENTAL MUSTARDS ONLY) IN THE PRAIRIE PROVINCES AND PEACE RIVER REGION OF BRITISH COLUMBIA; FOR SALE FOR USE IN CANOLA IN THE OKANAGAN AND CRESTON VALLEY AREAS OF BRITISH COLUMBIA, SOUTHERN ONTARIO, QUEBEC AND THE MARITIMES ONLY; FOR SALE FOR USE ON ORIENTAL MUSTARD (CONDIMENT AND OIL SEED TYPES) IN WESTERN CANADA ONLY. ALSO FOR SALE FOR USE ON SUNFLOWER, LAURENTIAN RUTABAGA AND BRASSICA CARINATA

AGRICULTURAL

READ THE LABEL AND THIS BOOKLET BEFORE USING

GUARANTEE: Ethametsulfuron-methyl 75%

Warning, contains the allergen sulfites.

REGISTRATION NO.: 23569 PEST CONTROL PRODUCTS ACT

DO NOT SELL WATER SOLUBLE BAGS SEPARATELY.

NET CONTENTS: 80-800 GRAMS
(1-10 x 80 gram water soluble bags)

E. I. DU PONT CANADA COMPANY
AGRICULTURAL PRODUCTS
P.O BOX 2300, STREETSVILLE
MISSISSAUGA, ONTARIO
L5M 2J4
1-800-667-3925

PRECAUTIONS:

- **KEEP OUT OF REACH OF CHILDREN.**
- Avoid breathing spray mist.
- Avoid contact with skin, eyes and clothing.
- Do not contaminate any body of water.
- Wear chemical resistant gloves during mixing, loading and cleanup.
- Wash hands and face before eating, drinking, smoking and using the toilet.
- Take a shower immediately after use.
- Wash protective clothing in detergent and hot water, separately from household laundry, before reuse.
- **DO NOT APPLY BY AIR.**
- When applied as a tank-mix combination, read and observe all label directions, including rates, restrictions, and grazing limitations for each product used in the tank-mix. Follow the more stringent label precautionary and PPE measures for mixing, loading and applying stated on both product labels.

- Wear long-sleeved shirt, long pants, socks and chemical-resistant gloves and footwear during mixing, loading, application, clean-up and repair. Chemical-resistant gloves are not required while operating groundboom sprayers.

FIRST AID:

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

For medical emergencies call 1-800-441-3637 (24 hours).

TOXICOLOGICAL INFORMATION: Treat Symptomatically.

ENVIRONMENTAL HAZARDS

Observe buffer zones specified under Directions for Use.

Injury to or loss of desirable trees or vegetation may result from failure to observe the following. Do not apply, drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, recreational grounds or similar areas. Prevent drift of spray to desirable plants. Do not contaminate any body of water, including irrigation water that may be used on other crops.

DO NOT APPLY BY AIR. APPLY WITH GROUND EQUIPMENT ONLY.

Carefully clean sprayer after use, as spray tank residue may damage crops other than canola (rapeseed), brown condiment mustard and oriental mustard (condiment and oilseed types).

FOR USE ONLY AS AN AGRICULTURAL HERBICIDE.

The manufacturer has not tested all possible surfactants for compatibility, efficacy, crop damage, or adverse effects. Therefore, DuPont does not recommend the use of other surfactants not listed on the label and is not responsible for any consequences of such use.

PRIOR TO USE, REFER TO "MINIMUM CROP ROTATION GUIDELINES" SECTION OF LABEL.

TOXIC to aquatic organisms and non target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE.

Runoff:

To reduce runoff from treated areas into aquatic habitats, consider the characteristics and conditions of the site before treatment. Site characteristics and conditions that may lead to runoff include, but are not limited to: heavy rainfall, moderate to steep slope, bare soil, poorly draining soil (e.g. soils that are compacted or fine textured such as clay).

Avoid application of this product when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

Leaching:

The use of this chemical may result in contamination of groundwater, particularly in areas where soil pH>7, are permeable (e.g. sandy soil) and/or the depth to the water table is shallow.

GENERAL INFORMATION

MUSTER® Toss-N-Go® Herbicide is recommended for use on spring planted canola (rapeseed) varieties, brown condiment mustard and oriental mustard (condiment and oilseed types) grown in Canada.

MUSTER® Toss-N-Go® Herbicide is a 75% active ingredient herbicide dry flowable dispersible granule contained in a water-soluble film.

MUSTER® Toss-N-Go® Herbicide is to be mixed in water and applied as a uniform broadcast spray with a recommended surfactant, for selective control or suppression of certain broadleaf weeds in canola (rapeseed) as well as brown condiment mustard and oriental mustard (condiment and oilseed types). It is noncorrosive, nonflammable, nonvolatile and does not require heated storage. Do not graze or feed crop to livestock within 60 days of treatment. Do not harvest within 60 days of treatment.

MUSTER® Toss-N-Go® Herbicide MUST be applied with a recommended surfactant early postemergence, to the main flush of actively growing broadleaf weeds. Registered surfactants are AG-SURF*, AGRAL* 90, CITOWETT* PLUS, CANPLUS* 411, HI-MIX®, SURE-MIX™ and MERGE*. Warm, moist growing conditions promote active weed growth and enhance the activity of MUSTER® Toss-N-Go® Herbicide by allowing maximum foliar uptake and contact activity.

Weeds hardened off by cold weather or drought stress may not be adequately controlled or suppressed and regrowth may occur. Regrowth may also occur if crop competitiveness is impaired by thin stands and/or reduced vigour.

MUSTER® Toss-N-Go® Herbicide may only be applied with ground spray equipment. For best results, ensure thorough spray coverage of target weeds. See "Directions for Use" section of this label for complete use details.

MUSTER® Toss-N-Go® Herbicide is an ALS inhibitor herbicide and thus rapidly inhibits the growth of susceptible weeds; however, typical symptoms (discolouration) of dying weeds may not be noticeable for 1 to 3 weeks after application depending upon growing conditions and weed susceptibility. Degree of control and duration of effect depend on weed sensitivity, weed size, growing conditions, and spray coverage.

DIRECTIONS FOR USE

- Do not re-enter treated fields until 12 hours after application.
- Apply only when the potential for drift to areas of human habitation or areas of human activity (houses, cottages, schools and recreational areas) is minimal. Take into consideration wind speed, wind direction temperature, application equipment and sprayer settings.

Field Sprayer Application:

- **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty.
- **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. The best available application strategies which minimize off-site drift, including meteorological conditions (e.g. wind direction, low wind speed) and spray equipment (e.g. coarse droplet sizes, minimizing height above canopy), should be used. Applicators must, however observe the specified buffer zones for protection of sensitive aquatic habitats.
- Boom height must be 60cm or less above the crop or ground.

Buffer Zones:

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, rangelands, riparian areas and shrub lands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks marshes, streams, reservoirs and wetlands) and marine/estuarine habitats.

Use of the following spray methods or equipment DO NOT require a buffer zone: hand-held or backpack sprayer and spot treatment.

Method of Application	Crop	Region of Production	Buffer Zones (metres) Required for the Protection of:		
			Aquatic Habitat of Depths:		Terrestrial Habitat
			Less than 1m	Greater than 1m	
Field sprayer*	Mustard (Condiment Brown and Oriental including both the Condiment types and the Oil seed types)	Prairie provinces and Peace River Region of British Columbia and Western Canada	3	2	40
	Rutabaga	National	3	1	30
	Canola	Southern Ontario, Québec and the Maritimes (New Brunswick, Prince Edward Island and Nova Scotia)	3	1	30
		Prairie Provinces, Peace River, Okanagan and Creston Valley areas of British Columbia	4	2	55
	Sunflower and <i>Brassica carinata</i>	National	4	2	55

*For field sprayer application, buffer zones can be reduced with the use of drift reducing spray shields. When using a spray boom fitted with a full shield (shroud, curtain) that extends to the crop canopy or ground, the labelled buffer zone can be reduced by 70%. When using a spray boom where individual nozzles are fitted with cone-shaped shields are no more than 30 cm above the crop canopy or ground, the labelled buffer zone can be reduced by 30%.

NOTE: Applicators may recalculate a site-specific buffer zone by combining information on current weather conditions and spray configuration for the following applications: all airblast applications, and for field and aerial applications which specify the following droplet size category wording on the product label: 'DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification.' To access the Buffer Zone Calculator, please visit the Pest Management Regulatory Agency web site.

When a tank mixture is used, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture.

MUSTER® Toss-N-Go® HERBICIDE USED ALONE:

SPRING CANOLA (RAPESEED)

MUSTER® Toss-N-Go® HERBICIDE + NON-IONIC SURFACTANT (AG-SURF*, AGRAL* 90 OR CITOWETT* PLUS)

Apply MUSTER® Toss-N-Go® Herbicide at 20 or 30 grams per hectare with a recommended surfactant at 2 litres surfactant per 1000 litres of spray volume (see "Mixing Instructions" section of label). For optimum weed control apply MUSTER® Toss-N-Go® Herbicide at the cotyledon to six (6) leaf stage of the target weeds, see specific recommendation for stinkweed. For best results apply to the main flush of young, actively growing weeds. Apply before the crop canopy prevents thorough coverage of the small target weeds. Thorough spray coverage of the small target weeds is important for good weed control. When very high weed seedling populations occur, larger weed seedlings may interfere with coverage of smaller seedlings, and control may be reduced. Weeds that occur after application may not be controlled. MUSTER® Toss-N-Go® Herbicide can be applied to canola (rapeseed) from the two (2) leaf stage to the beginning of bolting.

20 g/ha	WEEDS CONTROLLED Flixweed (spring seedlings) Green Smartweed Hemp-nettle Wild Mustard	WEEDS SUPPRESSED[†] Stinkweed
30 g/ha	WEEDS CONTROLLED Weeds controlled at 20 g/ha plus Stinkweed	WEEDS SUPPRESSED[†] Redroot Pigweed

[†] Weed suppression is a visual reduction in weed competition (reduced population or vigour) as compared to an untreated area. Degree of suppression will vary with size of weed and environmental conditions prior to and following treatment.

BROWN CONDIMENT MUSTARD AND ORIENTAL MUSTARD (CONDIMENT AND OILSEED TYPES)

MUSTER® Toss-N-Go® Herbicide + NON-IONIC SURFACTANT (AG-SURF*, AGRAL* 90 OR CITOWETT* PLUS)

Apply MUSTER® Toss-N-Go® Herbicide at 20 grams per hectare with a recommended surfactant at 2.0 litres per 1000 litres of spray volume (see "Mixing Instructions" section of label). Apply from the 4-leaf to late rosette stage of brown condiment mustard and oriental mustard (condiment and oilseed types), when target weeds are in the cotyledon to 6-leaf stage. For best results, apply to the main flush of young actively growing weeds. Thorough spray coverage of weeds is important for good control. When very high weed seedling population occurs, larger weeds may interfere with coverage of smaller seedlings, and control may be reduced.

- Do not use more than 20 grams per hectare on brown condiment mustard and oriental mustard (condiment and oilseed types) or injury may result.
- Do not use on yellow mustard varieties.
- Do not apply in combination with other pesticides.

20 g/ha	WEEDS CONTROLLED Flixweed (spring seedlings) Green Smartweed Hemp-nettle Wild Mustard	WEEDS SUPPRESSED[†] Stinkweed
----------------	--	--

[†] Weed suppression is a visual reduction in weed competition (reduced population or vigour) as compared to an untreated area. Degree of suppression will vary with size of weed and environmental conditions prior to and following treatment.

NOTE:

If rain occurs soon after application control may be reduced. At least 4 - 6 hours of dry weather are needed to allow MUSTER® Toss-N-Go® Herbicide to be absorbed by weed foliage. Environmental conditions that slow the drying of MUSTER® Toss-N-Go® Herbicide on the foliage such as high relative humidity, cool air temperature or cloud cover, may increase the time required.

SPECIFIC WEED RECOMMENDATIONS:

Stinkweed: For best results apply MUSTER® Toss-N-Go® Herbicide plus recommended surfactant to actively growing emerged stinkweed that are in the 1-4 leaf stage. Ensure thorough coverage.

FOR CONTROL OF WILD MUSTARD IN SPRING-SEEDED CANOLA IN SOUTHERN ONTARIO:

MUSTER® Toss-N-Go® HERBICIDE + NON-IONIC SURFACTANT (AG-SURF*, AGRAL* 90 OR CITOWETT* PLUS)

Apply MUSTER® Toss-N-Go® Herbicide at 15 grams per hectare with a recommended surfactant at 2 litres surfactant per 1000 litres of spray volume. This rate should be used only in Southern Ontario. Apply when wild mustard is in the cotyledon to 6-leaf stage, before the crop begins to bolt. Land treated with MUSTER® Toss-N-Go® Herbicide must not be planted to crops other than winter wheat in the year of treatment; soybeans or spring cereals in the year following treatment.

MUSTER® Toss-N-Go® HERBICIDE TANK MIXTURES:

TANK MIXTURES IN SPRING CANOLA (PRAIRIE PROVINCES AND PEACE RIVER REGION OF BRITISH COLUMBIA)

MUSTER® Toss-N-Go® Herbicide Plus POAST*Herbicide

For control of broadleaf weeds and grasses in canola (rapeseed) apply a tank mixture of MUSTER® Toss-N-Go® Herbicide plus POAST* Herbicide. Use 20 or 30 grams per hectare MUSTER® Toss-N-Go® Herbicide plus 0.8 to 1.9 litres per hectare POAST* Herbicide plus the recommended amount of MERGE*

adjuvant as stated on the POAST* Herbicide label. Consult the POAST* Herbicide label for restrictions and desired use rates. Additional surfactant is not required when using MERGE*. For best results follow the timing restrictions and specific weed control recommendations stated on each label. Stinkweed will be controlled by 20 grams of MUSTER® Toss-N-Go® Herbicide per hectare when used in this mixture with POAST* Herbicide and MERGE*.

MUSTER® Toss-N-Go® Herbicide Plus ASSURE® II Herbicide

For control of broadleaf weeds and grasses in canola (rapeseed), apply a tank mixture of MUSTER® Toss-N-Go® Herbicide plus ASSURE® II Herbicide. Use MUSTER® Toss-N-Go® Herbicide at 20 or 30 grams per hectare plus 0.38 - 0.5 litre per hectare ASSURE® II Herbicide plus CANPLUS* 411 or MERGE* at 5-10 litres per 1000 litres of spray solution or SURE-MIX™ at 5 litres per 1000 litres of spray solution. Additional surfactant is not required when using CANPLUS* 411, HI-MIX®, MERGE* or SURE-MIX™. For best results follow the timing restrictions and specific weed control recommendations stated on each label. Consult the ASSURE® II Herbicide label for desired use rates, grass weeds controlled and additional restrictions regarding the use of ASSURE® II Herbicide. MUSTER® Toss-N-Go® Herbicide will control stinkweed at the 20 grams per hectare rate when used with HI-MIX®, MERGE* or SURE-MIX™ adjuvant and ASSURE® II Herbicide.

TANK MIXTURE IN ORIENTAL MUSTARD (CONDIMENT AND OILSEED TYPES) FOR WESTERN CANADA:

MUSTER® Toss-N-Go® Herbicide Plus ASSURE® II Herbicide

For control of broadleaf weeds and grasses in oriental mustard (condiment and oilseed types), apply a tank mixture of MUSTER® Toss-N-Go® Herbicide plus ASSURE® II Herbicide. Use MUSTER® Toss-N-Go® Herbicide at 20 grams per hectare plus 0.38 - 0.5 litre per hectare ASSURE® II Herbicide plus MERGE* at 5-10 litres per 1000 litres of spray solution or SURE-MIX™ at 5 litres per 1000 litres of spray solution. Additional surfactant is not required when using MERGE* or SURE-MIX™. For best results follow the timing restrictions and specific weed control recommendations stated on each label. Consult the ASSURE® II Herbicide label for desired use rates, grass weeds controlled and additional restrictions regarding the use of ASSURE® II Herbicide. MUSTER® Toss-N-Go® Herbicide will control stinkweed at the 20 grams per hectare rate when used with MERGE* or SURE-MIX™ adjuvant and ASSURE® II Herbicide.

- Do not use on yellow mustard varieties.

When a tank mixture is used, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture.

MIXING INSTRUCTIONS:

Do not mix, load or apply within 15 metres of all wells, including abandoned wells and drainage wells.

1. Fill the tank with clean water with agitator running. Use a minimum of 100 litres of spray volume per hectare.
2. Add the required amount of MUSTER® Toss-N-Go® Herbicide with the agitator running. Continuous agitation is required for a uniform suspension and application.
3. Once the MUSTER® Toss-N-Go® Herbicide is completely in suspension, add a recommended non-ionic surfactant at 2 litres per 1000 litres of spray solution. Antifoaming agents may be needed.
4. FOR TANK MIXTURES WITH ASSURE® II Herbicide OR POAST* Herbicide: If tank mixing MUSTER® Toss-N-Go® Herbicide with ASSURE® II Herbicide or POAST* Herbicide, make sure the MUSTER® Toss-N-Go® Herbicide bags have completely dissolved, then add the required amount of ASSURE® II Herbicide or POAST* Herbicide to the tank with the agitator running. Once the

herbicide is completely in suspension, SLOWLY add the required amount of partner surfactant: ASSURE® II Herbicide with CANPLUS* 411, MERGE* or SURE-MIX™, POAST* Herbicide with MERGE*. ADDITIONAL SURFACTANT IS NOT REQUIRED.

On repeat tank loads, ensure that the amount of spray solution left in the tank from the previous load is less than 10% of the volume about to be sprayed.

Use the spray preparation of MUSTER® Toss-N-Go® Herbicide within 24 hours or product degradation may occur. If spray preparation is left standing without agitation, thoroughly agitate and ensure product is resuspended before resuming spray operation.

Avoid over-filling of spraytank. Mix only enough product for the job at hand.

During spraying operation, make scheduled checks of spray equipment to help ensure proper application.

NOTE:

Do not mix the soluble bags with liquid fertilizers, substances that contain boron or substances that release free chlorine. Mixing the water soluble bags with any of these compounds will result in an insoluble substance in the tank.

- Drop intact water-soluble bags (WSB) into the spray tank. The WSB and pesticide will dissolve readily in water.
- Do not allow the WSB to become wet prior to use. Do not handle individual WSB with wet hands or wet gloves as this may cause breakage. Do not remove WSB from over-wrap container except for immediate use.
- Do not open WSB or attempt to remove product from them. The contents are not designed to be re-measured or subdivided.
- Do not open or use punctured WSB.
- If broken WSB is found when the over-wrap bag is opened, avoid contact with, and inhalation of the product. Handle with chemical-resistant gloves and dispose of broken WSB according to the DISPOSAL section.

In some cases, tank mixing a pest control product with another pest control product or a fertilizer can result in biological effects that could include, but are not limited to: reduced pest efficacy or increased host crop injury. The user should contact E.I du Pont Canada Company at 1-800-667-3925 for information before mixing any pesticide or fertilizer that is not specifically recommended on this label. The user assumes the risk of losses that result from the use of tank mixes that do not appear on this label or that are not specifically recommended by E.I du Pont Canada Company.

APPLICATION INSTRUCTIONS:

Apply the spray mixture with ground equipment only, using a minimum of 100 litres spray volume per hectare. Only use a properly calibrated sprayer that will ensure thorough coverage of target weeds and give a uniform spray pattern. When calibrating a sprayer, always use clean water and calibrate away from well sites or any sources of water. Flat fan nozzles are recommended. Use 50-mesh (metal or nylon) screens or larger. A spray pressure of 210-275 kPa is recommended.

Continuous agitation is required to keep MUSTER® Toss-N-Go® Herbicide in suspension. Avoid overlapping, and shut off spray booms while starting, turning, slowing or stopping to prevent over application.

NOTE: Care must be taken to prevent drift to desirable plants or non-target agricultural land.

SPRAYER CLEANUP:

On completion of spraying with MUSTER® Toss-N-Go® Herbicide, thoroughly remove all traces of herbicide from mixing and spray equipment in order to avoid subsequent injury to sensitive crops other than canola (rapeseed).

Follow these instructions:

1. Drain tank and flush tank, boom and hoses with clean water for a minimum of ten minutes. Visually inspect tank to assure removal of all visible residues of MUSTER® Toss-N-Go® Herbicide. If necessary, repeat step 1. **DO NOT CLEAN SPRAYER NEAR WELL OR WATER SOURCE OR NEAR TO DESIRABLE VEGETATION.**
2. Fill tank with clean water while adding 1 litre household ammonia (containing a minimum 3% ammonia) per 100 litres of water. Flush solution through boom and hoses, and then add more water to completely fill tank. Allow to sit for 15 minutes with agitation. Again flush the hoses, boom and nozzles with the cleaning solution and drain tank.
3. Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.
4. Repeat Step 2.
5. Thoroughly rinse the tank with clean water for a minimum of 5 minutes flushing water through the hoses and boom.

CAUTION:

DO NOT use ammonia with chlorine bleach. Using ammonia with chlorine bleach will release a gas with a musty chlorine odour which may cause eye, nose, throat, and lung irritation. DO NOT clean equipment in an enclosed area.

MINIMUM CROP ROTATION GUIDELINES:

Minimum interval is that time from the last application of MUSTER® Toss-N-Go® Herbicide to date of planting the rotational crop.

**INTERVAL PRIOR TO PLANTING
(MONTHS AFTER APPLICATION)**

10 months	Spring Wheat, Durum Wheat, Barley, Oats, Flax
22 months	Canola, Lentils, Peas, Fababeans, Tame Mustard, Alfalfa, Canary Grass, Dry Beans, Fescue, Red Clover

All other crops field bioassay at 22 months.

NOTE:

Wherever MUSTER® Toss-N-Go® Herbicide is used on land previously treated with E. I. du Pont Canada Company GLEAN® Herbicide or ALLY® Herbicide, read the rotational guidelines on both labels and follow the label with the longest interval stated for your situation.

FIELD BIOASSAY:

Select a representative area or areas of the field previously treated with MUSTER® Toss-N-Go® Herbicide to plant your bioassay crop(s). Be sure to consider factors such as size of field, soil texture, drainage and turn-around areas when selecting the site(s) that are most representative of the soil conditions in the field. On large fields, more than one site may be needed in order to obtain reliable results. Plant the test strips perpendicular to the direction in which the field was sprayed. The strips should be long enough to cross the width of several spray swaths. Large test strip areas are more reliable than small ones.

Use standard tillage and seeding equipment to plant the bioassay. Prepare a seed bed and plant the crops and varieties you want the option of growing the following year. It is important to use the same planting time, conditions, techniques and cultural practices you normally use to plant and grow the bioassay crop(s). Also plant into an adjacent area not treated with MUSTER® Toss-N-Go® Herbicide to use as a comparison.

As the crop(s) emerges and grows, examine these key points in MUSTER® Toss-N-Go® Herbicide treated and non-treated areas:

Crop stand	Root development	Rate of growth
Plant colour and vigour	Yield	

Allow the bioassay crop(s) to grow to maturity while making your observations. Do not overspray the test strips with herbicides that may damage the bioassay crop(s). If the bioassay indicates that MUSTER® Toss-N-Go® Herbicide residues are still present, continue cropping only to those crops listed on the label and do not rotate to other crops until bioassay results indicate that susceptible crops are growing normally.

USE PRECAUTIONS:

- DO NOT apply this product directly to freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs, ditches and wetlands), estuaries or marine habitats.
- DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.
- DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty.
- Do not use more than 20 grams per hectare per year on brown condiment mustard or oriental mustard (condiment and oilseed types).
- Do not use more than 30 grams per hectare per year on canola (rapeseed).
- Avoid overspray or drift to important wildlife habitats.
- Licensed canola (rapeseed) varieties, brown condiment mustard and oriental mustard (condiment and oilseed types) have shown acceptable tolerance to MUSTER® Toss-N-Go® Herbicide. Consult the E. I. du Pont Canada Company prior to use on unlicensed varieties for crop tolerance information.
- Do not apply to irrigated land where tail water will be used to irrigate other crop land.
- Do not contaminate irrigation water.
- Do not use on highly variable soils that have large gravelly or sandy areas, eroded knolls, or calcium deposits.
- Do not apply MUSTER® Toss-N-Go® Herbicide to canola (rapeseed), brown condiment mustard or oriental mustard (condiment and oilseed types) or *Brassica carinata* that are stressed by severe conditions such as drought, low fertility, saline soils, waterlogged soils (soils at or near field capacity), disease or insect damage as crop injury may result. Drought, disease or insect damage following application may also result in crop injury.
- Do not apply to canola (rapeseed), brown condiment mustard or oriental mustard (condiment and oilseed types) or *Brassica carinata* undersown with legumes or grasses as injury to the forage may result.
- Do not remove soil from treated fields for use on lawns, in gardens or for backfill.

- Heavy rainfall soon after application may result in visual crop injury or possible yield reduction. Conditions such as thin crop stand, application prior to the 2-leaf stage of canola or 4-leaf stage of brown condiment mustard and oriental mustard (condiment and oilseed types), sandy soil or low soil organic matter may increase the severity of injury.
- The additive effect of soil residues from the use of MUSTER® Toss-N-Go® Herbicide and ASSERT* Herbicide on the same land has not been determined. Crop rotation guidelines and minimum rotation intervals are not known and injury to rotational crops other than wheat (excluding durum) may occur. Wherever land will be treated with MUSTER® Toss-N-Go® Herbicide and has been or will be treated with ASSERT* Herbicide, plant only wheat (excluding durum) after both herbicides have been used, until a bioassay (see "Field Bioassay" section of this label) demonstrates that other crops can be successfully grown.

NOTE TO BUYER/USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR SPECIAL USE APPLICATIONS:

The DIRECTIONS FOR USE for this product for the use described below were developed by persons other than the E. I. du Pont Canada Company. and accepted for registration by Health Canada under the User Requested Minor Use Label Expansion program. The E. I. du Pont Canada Company. itself makes no representation or warranty with respect to product performance (efficacy) and crop tolerance (phytotoxicity) of this product when used on the crops below. Accordingly, the Buyer and User assume all liability arising, and agree to hold The E. I. du Pont Canada Company. harmless from any claims based on efficacy or phytotoxicity in connection with this use.

FOR USE ON SPRING SEEDED CANOLA (Rapeseed) VARIETIES IN QUEBEC AND IN THE MARITIMES (NEW BRUNSWICK, PRINCE EDWARD ISLAND AND NOVA SCOTIA):

For postemergent control of wild mustard in canola, apply MUSTER® Toss-N-Go® Herbicide at 15 grams per hectare with a recommended NON-IONIC SURFACTANT (AG-SURF*, AGRAL* 90, or CITOWETT* PLUS) at 2 litres per 1000 litres of spray volume OR a recommended adjuvant (SURE-MIX™) at 5 litres per 1000 litres of spray volume.

For optimum weed control apply MUSTER® Toss-N-Go® Herbicide at the cotyledon to six (6) leaf stage of wild mustard. For best results apply to the main flush of young, actively growing weeds. Apply before the crop canopy prevents thorough coverage of the small target weeds. Wild mustard seedlings that occur after application may not be controlled. MUSTER® Toss-N-Go® Herbicide can be applied to canola (rapeseed) from the two (2) leaf stage to the beginning of bolting.

Land treated with MUSTER® Toss-N-Go® Herbicide must not be planted to crops other than spring wheat, barley or oats with a recropping interval of 10 months and winter wheat, which may be planted in the year of treatment.

Apply with ground equipment only. Make only one application per growing season. Do not harvest crop within 60 days of treatment.

Refer to other sections of this booklet for additional application and/or use precaution instructions.

SPRING SEEDED CANOLA (Rapeseed) VARIETIES IN THE OKANAGAN AND CRESTON VALLEY AREAS OF BRITISH COLUMBIA:

MUSTER® Toss-N-Go® Herbicide + Non-Ionic Surfactant (AG-SURF®, AGRAL® 90 OR CITOWETT® PLUS)

Apply MUSTER® Toss-N-Go® Herbicide at 20 or 30 grams per hectare with a recommended surfactant (AG-SURF, AGRAL 90 OR CITOWETT PLUS) at 2 litres surfactant per 1000 litres of spray volume (see "Mixing Instructions" section of label).

For optimum weed control apply MUSTER® Toss-N-Go® Herbicide at the cotyledon to six (6) leaf stage of the target weeds, see specific recommendation for stinkweed. For best results apply to the main flush of young, actively growing weeds. Apply before the crop canopy prevents thorough coverage of the small target weeds. Thorough spray coverage of the small target weeds is important for good weed control. When very high weed seedling populations occur, larger weed seedlings may interfere with coverage of smaller seedlings, and control may be reduced. Weeds that occur after application may not be controlled. MUSTER® Toss-N-Go® Herbicide can be applied to canola (rapeseed) from the two (2) leaf stage to the beginning of bolting.

20 g/ha	WEEDS CONTROLLED Flixweed (spring seedlings) Green Smartweed Hemp-nettle Wild Mustard	WEEDS SUPPRESSED* Stinkweed
30 g/ha	WEEDS CONTROLLED Weeds controlled at 20 g/ha plus Stinkweed	WEEDS SUPPRESSED* Redroot Pigweed

*Weed suppression is a visual reduction in weed competition (reduced population or vigour) as compared to an untreated area. Degree of suppression will vary with size of weed and environmental conditions prior to and following treatment.

Apply with ground equipment only. Make only one application per growing season. Do not harvest crop within 60 days of treatment.

Refer to other sections of this booklet for additional application and/or use precaution instructions.

FOR USE ON SUNFLOWER:

For postemergent control of weeds listed on the "MUSTER® Toss-N-Go® Herbicide" section of the label, apply MUSTER® Toss-N-Go® Herbicide at 20-30 grams per hectare with a recommended NON-IONIC SURFACTANT (AG-SURF®, AGRAL® 90, or CITOWETT® PLUS) at 2 litres per 1000 litres of spray volume. Use a minimum spray volume of 100 L/ha.

For optimum weed control apply MUSTER® Toss-N-Go® Herbicide at the cotyledon to six (6) leaf stage of target weeds, see specific recommendations for stinkweed. For best results apply to the main flush of young, actively growing weeds. Apply before the crop canopy prevents thorough coverage of the small target weeds. Weed seedlings that occur after application may not be controlled. MUSTER® Toss-N-Go® Herbicide can be applied to sunflower from 15 to 45 cm in height (2 to 8 leaf stage).

Do not graze the treated crops or cut for hay; sufficient data are not available to support such use.

Land treated with MUSTER® Toss-N-Go® Herbicide must not be planted to crops other than spring wheat, barley or oats with a recropping interval of 10 months. See the "INTERVAL PRIOR TO PLANTING" section of the label for additional recropping restrictions and recommendations.

Apply with ground equipment only. Make only one application per growing season. Do not harvest crop within 60 days of treatment.

Refer to other sections of this booklet for additional application and/or use precaution instructions.

FOR USE ON LAURENTIAN RUTABAGA:

For postemergent control of wild mustard, apply MUSTER® Toss-N-Go® Herbicide at 15 grams per hectare with a recommended NON-IONIC SURFACTANT (AG-SURF®, AGRAL® 90, or CITOWETT® PLUS) at 2 litres per 1000 litres of spray volume OR a recommended adjuvant (SURE-MIX™) at 5 litres per 1000 litres of spray volume.

Apply MUSTER® Toss-N-Go® Herbicide from the cotyledon to the 6-leaf stage of wild mustard. For best results apply to the main flush of young, actively growing weeds. Apply before the crop canopy prevents thorough coverage of the small target weeds. Weeds that emerge after application may not be controlled.

MUSTER® Toss-N-Go® Herbicide can be applied post-emergent to Laurentian rutabaga prior to crop bolting.

Apply with ground equipment only. Use a spray volume of at least 100 L/Ha. Make only one application per growing season. Do not harvest crop within 45 days of treatment.

See the "INTERVAL PRIOR TO PLANTING" section of the label for additional recropping restrictions and recommendations.

Refer to other sections of this booklet for additional application and/or use precaution instructions.

FOR USE ON BRASSICA CARINATA**MUSTER® Toss-N-Go® Herbicide + NON-IONIC SURFACTANT (AG-SURF*, AGRAL* 90 OR CITOWETT* PLUS)**

Apply MUSTER® Toss-N-Go® Herbicide at 20 grams per hectare with a recommended surfactant at 2.0 litres per 1000 litres of spray volume (see "Mixing Instructions" section of label). Apply from the 4-leaf to late rosette stage of Brassica Carinata, when target weeds are in the cotyledon to 6-leaf stage. For best results, apply to the main flush of young actively growing weeds. Thorough spray coverage of weeds is important for good control. When very high weed seedling population occurs, larger weeds may interfere with coverage of smaller seedlings, and control may be reduced.

- Do not use more than 20 grams per hectare on Brassica Carinata or injury may result.

20 g/ha	WEEDS CONTROLLED	WEEDS SUPPRESSED [†]
	Flixweed (spring seedlings)	Stinkweed
	Green Smartweed	
	Hemp-nettle	
	Wild Mustard	

[†] Weed suppression is a visual reduction in weed competition (reduced population or vigour) as compared to an untreated area. Degree of suppression will vary with size of weed and environmental conditions prior to and following treatment.

NOTE:

If rain occurs soon after application control may be reduced. At least 4 - 6 hours of dry weather are needed to allow MUSTER® Toss-N-Go® Herbicide to be absorbed by weed foliage. Environmental conditions that slow the drying of MUSTER® Toss-N-Go® Herbicide on the foliage such as high relative humidity, cool air temperature or cloud cover, may increase the time required.

SPECIFIC WEED RECOMMENDATIONS:

Stinkweed: For best results apply MUSTER® Toss-N-Go® Herbicide plus recommended surfactant to actively growing emerged stinkweed that are in the 1-4 leaf stage. Ensure thorough coverage.

Refer to other sections of this booklet for additional application and/or use precaution instructions.

RESISTANCE-MANAGEMENT RECOMMENDATIONS:

For resistance management, MUSTER® Toss-N-Go® Herbicide is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to MUSTER® Toss-N-Go® Herbicide and other Group 2 herbicides. The resistance biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- ◆ Where possible rotate the use of MUSTER® Toss-N-Go® Herbicide or other Group 2 herbicides with different herbicide groups that control the same weeds in a field.
- ◆ Use tank mixtures with herbicides from a different group when such use is permitted
- ◆ Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural, biological and other chemical control practices.
- ◆ Monitor treated weed populations for resistance development.
- ◆ Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed.
- ◆ Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- ◆ For further information or to report suspected resistance, contact your local E. I. du Pont Canada Company representative or the E. I. du Pont Canada Company hotline at 1-800-667-3925 for further information.

STORAGE:

Store product in original container only, away from fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed. Keep water soluble bags in original outer bag away from moisture.

DISPOSAL:

1. Make the empty container unsuitable for further use.
2. Dispose of the container in accordance with provincial requirements.
3. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER:

This control product is to be used only in accordance with the directions on this label. It is an offence under the PEST CONTROL PRODUCTS ACT to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

The seller warrants that the purchase by the buyer and the use of this product, as such, will not infringe any Canadian patent.

®/™ Registered trademarks/trademarks of E.I. du Pont de Nemours and Company. E. I. du Pont Canada Company is a licensee. Member of CropLife.

* All other products mentioned are trademarks of their respective companies.

<<Water Soluble Bag Label>>

MUSTER® Toss-N-Go® HERBICIDE

KEEP OUT OF REACH OF CHILDREN

SEE DIRECTIONS FOR USE BROCHURE FOR COMPLETE PRECAUTIONARY STATEMENTS

GUARANTEE: ETHAMETSULFURON METHYL 75%

DRY FLOWABLE

Warning, contains the allergen sulfites.

NET CONTENTS: 80 GRAMS