

<b>GROUP</b>	<b>28</b>	<b>INSECTICIDE</b>
--------------	-----------	--------------------

**DUPONT™ CORAGEN™ insecticide**

SUSPENSION

FOR SALE FOR USE ON ROOT AND TUBER VEGETABLES GROUP, FRUITING VEGETABLES GROUP, BRASSICA VEGETABLES GROUP, LEAFY VEGETABLES GROUP, LEGUME VEGETABLES GROUP CUCURBIT VEGETABLES GROUP, CORN (FIELD, POP, SWEET, AND SEED), GRASS FORAGE, FODDER, AND HAY GROUP, NON-GRASS ANIMAL FEEDS GROUP, MINT, OKRA, PEANUTS, GREEN ONION SUB GROUP, GLOBE ARTICHOKE, HOPS, GREENHOUSE CUCUMBERS, GREENHOUSE TOMATOES, GREENHOUSE EGGPLANT, GREENHOUSE PEPPERS, OILSEEDS GROUP AND CEREALS

COMMERCIAL

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

Contains 1,2-benzisothiazolin-3-one at 0.019% as a preservative  
or

Contains 5-chloro-2-methyl-4-isothiazolin-3-one at 0.00084% and 2-methyl-4-isothiazolin-3-one at 0.00028% and 2-bromo-2-nitropropane-1,3-diol at 0.0088% as preservatives

**GUARANTEE:** Chlorantraniliprole 200 g/L

REGISTRATION NO. 28982 PEST CONTROL PRODUCTS ACT

**NET CONTENTS:** 0.5, 3.79, 100 LITRES

E.I. DU PONT CANADA COMPANY  
AGRICULTURAL PRODUCTS  
P.O. BOX 2300, STREETSVILLE  
MISSISSAUGA, ONTARIO  
L5M 2J4  
1-800-667-3925

**PRECAUTIONS:**

- **KEEP OUT OF REACH OF CHILDREN.**
- Wear a long sleeved shirt, long pants and chemical resistant gloves while mixing /loading, applying and during clean up and repair.
- Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables are available, use detergent and hot water. Keep and wash PPE separately from other laundry.
- Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- For agricultural uses, do not enter or allow worker entry into treated areas during the restricted entry interval of 12 hours.
- See the Aerial Application Label Instructions section of this label for additional precautions specific to aerial application.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's website at [www.croplife.ca](http://www.croplife.ca).

**ENVIRONMENTAL HAZARDS:**

- Toxic to aquatic organisms. Observe buffer zones specified under **DIRECTIONS FOR USE.**
- Toxic to certain beneficial insects. Minimize spray drift to reduce harmful effects on beneficial insects in habitats next to the application site such as hedgerows and woodland.
- Chlorantraniliprole is persistent and may carryover. It is recommended that any products containing chlorantraniliprole not be used in areas treated with this product during the previous season.
- The use of this chemical may result in contamination of groundwater particularly in areas where soil is permeable (e.g. sandy soil) and/or the depth to the water table is shallow.
- To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.
- Avoid application when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.
- **DO NOT** apply this product directly to freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs, and wetlands), estuarine/marine habitats.
- **DO NOT** contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

**STORAGE:**

Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed. To prevent contamination, store this product away from food or feed.

**DISPOSAL:**

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

**FIRST AID:**

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**IF SWALLOWED:** Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

For medical emergencies call 1-800-441-3637 (24 hours).

**TOXICOLOGICAL INFORMATION:** Treat symptomatically. This product contains a petroleum distillate. Vomiting may cause aspiration pneumonia.

**NOTICE TO USER:**

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

**DUPONT™ CORAGEN™ insecticide**

SUSPENSION

FOR SALE FOR USE ON ROOT AND TUBER VEGETABLES GROUP, FRUITING VEGETABLES GROUP, BRASSICA VEGETABLES GROUP, LEAFY VEGETABLES GROUP, LEGUME VEGETABLES GROUP, CUCURBIT VEGETABLES GROUP, CORN (FIELD, POP, SWEET, AND SEED), GRASS FORAGE, FODDER, AND HAY GROUP, NON-GRASS ANIMAL FEEDS GROUP, MINT, OKRA, PEANUTS, GREEN ONION SUB GROUP, GLOBE ARTICHOKE, HOPS, GREENHOUSE CUCUMBERS, GREENHOUSE TOMATOES, GREENHOUSE EGGPLANT, GREENHOUSE PEPPERS, OILSEEDS GROUP AND CEREALS

COMMERCIAL

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

Contains 1,2-benzisothiazolin-3-one at 0.019% as a preservative

or

Contains 5-chloro-2-methyl-4-isothiazolin-3-one at 0.00084% and 2-methyl-4-isothiazolin-3-one at 0.00028% and 2-bromo-2-nitropropane-1,3-diol at 0.0088% as preservatives

**GUARANTEE:** Chlorantraniliprole 200 g/L

REGISTRATION NO. 28982 PEST CONTROL PRODUCTS ACT

**NET CONTENTS:** 0.5, 3.79, 100 LITRES

E.I. DU PONT CANADA COMPANY  
AGRICULTURAL PRODUCTS  
P.O. BOX 2300, STREETSVILLE  
MISSISSAUGA, ONTARIO  
L5M 2J4  
1-800-667-3925

## GENERAL INFORMATION

DUPONT™ CORAGEN™ insecticide is a liquid formulation that can be applied as a foliar spray, using ground application or aerial application (potatoes, legume vegetables, corn, oilseeds and cereals only) to control many important insect pests. DUPONT™ CORAGEN™ insecticide is mixed with water for application.

Apply at the recommended rates when insect populations reach locally determined economic thresholds. Consult your local extension specialist or certified crop advisor to determine appropriate threshold levels for treatment in your area.

Use sufficient water to obtain thorough, uniform coverage.

## DIRECTIONS FOR USE

**Do not allow effluent or runoff from greenhouses containing this product to enter lakes, streams, ponds or other waters.**

### ROOT AND TUBER VEGETABLES – Crop Group 1

**Spray Volume for Tuberos and Corm Vegetables:** Apply in a minimum finished spray volume of 100 L/ha by ground. For potatoes only, apply in a minimum finished spray volume of 50 L/ha by air.

SITE	PEST	RATE DUPONT™ CORAGEN™ insecticide /ha	SPECIFIC DIRECTIONS
POTATOES	Colorado potato beetle	250 mL to 500 mL	For potatoes, apply by either ground or aerial application equipment.
	European corn borer	250 mL to 375 mL	
ROOT AND TUBER VEGETABLES  Arracacha; Arrowroot; Artichoke, Chinese; Artichoke, Jerusalem; Beet, garden; Beet, sugar; Burdock, edible; Canna, edible; Carrot; Cassava, bitter and sweet; Celeriac (celery root); Chayote (root); Chervil, turnip-rooted; Chicory; Chufa; Dasheen (taro); Ginger; Ginseng; Horseradish; Leren; Parsley, turnip-rooted; Parsnip;	Diamondback moth	250 mL	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control.
	Cabbage looper		
	Black cutworm		
	Imported cabbageworm	250 mL to 375 mL	For Colorado potato beetle, use the higher application rate under heavy pest pressure and/or when larger larvae are present.
	Swede midge		
	Corn earworm		
European corn borer			
	Tobacco hornworm	Colorado potato beetle resistance management: DO NOT apply DuPont™ CORAGEN™ insecticide for Colorado potato beetle control if any Group 28 Insecticide	
	Tomato hornworm		
	Armyworm		
	Variegated		

Potato; Radish; Radish, oriental (daikon); Rutabaga; Salsify; Salsify, black; Salsify, Spanish; Skirret; Sweet potato; Tanier (cocoyam); Turmeric; Turnip; Yam bean; Yam, true	cutworm  Fall armyworm  Beet armyworm  Leafminers: Liriomyza sativae Liriomyza trifolii		<p>was used at planting as an in-furrow, soil or seed-piece treatment.</p> <p>For control of the European corn borer, time the application to coincide with peak egg hatch. Scout for European corn borer by monitoring egg laying and egg hatch to determine application timing.</p> <p>For black cutworm control, apply to foliage when rain is not expected in the next 24 hours. For optimal control, apply to smaller plants or when lower portions of the plant can receive adequate coverage.</p>
---	---	--	--

**RESTRICTIONS AND PRECAUTIONS: Tuberous and Corm Vegetables**

- For pests with a range of application rates, use the high rate under heavy pest pressure.
- Do not make more than 4 applications per season.
- Do not apply more than once every 3 days.
- Do not apply less than 1 day before harvest. Observe a 1-day PHI.
- Do not exceed a total of 1.125 litres DUPONT™ CORAGEN™ insecticide per ha per season.
- **Do not make a foliar application of DUPONT™ CORAGEN™ insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.**

## LEAFY VEGETABLES – Crop Group 4

**Spray Volume for Leafy Vegetables:** Apply in a minimum finished spray volume of 100 L/ha by ground.

SITE	PEST	RATE DUPONT™ CORAGEN™ insecticide /ha	SPECIFIC DIRECTIONS
LEAFY VEGETABLES Amaranth leafy, Arugula, Cardoon, Celery, Celery (Chinese), Celtuce, Chevril, Chrysanthemum (edible leaved), Chrysanthemum (garland), Corn Salad, Cress (garland), Cress (upland), Dandelion leaves, Dock, Endive, Florence Fennel, Lettuce (head & leaf), Orach, Parsley leaves, Purslane (garden), Purslane (winter), Radicchio, Rhubarb, Spinach, Spinach (vine), Spinach (New Zealand), Swiss Chard	Cabbage looper  Black cutworm	250 mL	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control.  For black cutworm control, apply to foliage when rain is not expected in the next 24 hours. For optimal control, apply to smaller plants or when lower portions of the plant can receive adequate coverage.
	Armyworm  Fall armyworm  Corn earworm  Beet armyworm  Leafminers: <i>Liriomyza sativae</i> <i>Liriomyza trifolii</i>	250 mL to 375 mL	

### RESTRICTIONS AND PRECAUTIONS: Leafy Vegetables

- Do not make more than 4 applications per season.
- Do not apply more than once every 3 days.
- Do not apply less than 1 day before harvest. Observe a 1-day PHI.
- Do not exceed a total of 1 litre of DUPONT™ CORAGEN™ insecticide per ha per season.
- **Do not make a foliar application of DUPONT™ CORAGEN™ insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.**

## BRASSICA VEGETABLES - Crop Group 5

**Spray Volume for Brassica Vegetables:** Apply in a minimum finished spray volume of 100 L/ha by ground.

SITE	PEST	RATE DUPONT™ CORAGEN™ insecticide / ha	SPECIFIC DIRECTIONS
BRASSICA VEGETABLES Broccoli, Broccoli (Chinese), Broccoli raab, Brussels Sprouts, Cabbage, Cabbage (Chinese, bok choy), Cabbage (Chinese, napa), Cabbage (Chinese mustard, choy), Cauliflower, Cavalo broccolo, Collards, Kale, Kohlrabi, Mizuna, Mustard Greens, Mustard Spinach, Rape Greens	Imported cabbage worm	250 ml	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control.  For optimum control, apply Hasten NT Spray Adjuvant* at an application rate of 0.25% v/v or MSO Contentrate with Leci- Tech* at an application rate of 0.5% v/v.  For black cutworm control, apply to foliage when rain is not expected in the next 24 hours. For optimal control, apply to smaller plants or when lower portions of the plant can receive adequate coverage.
	Diamondback moth		
	Cabbage looper		
	Black cutworm		
	Armyworm	250 ml to 375 ml	
	Fall armyworm		
	Beet armyworm		
	Corn earworm		
	Leafminers: <i>Liriomyza sativae</i> <i>Liriomyza trifolii</i>		

### RESTRICTIONS AND PRECAUTIONS: Brassica Vegetables

- Do not make more than 4 applications per season.
- Do not apply more than once every 3 days.
- Do not apply less than 3 days before harvest. Observe a 3-day PHI.
- Do not exceed a total of 1 litre of DUPONT™ CORAGEN™ insecticide per ha per season.
- **Do not make a foliar application of DUPONT™ CORAGEN™ insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.**



**LEGUME VEGETABLES - Crop Group 6**

**Spray Volume for Legume Vegetables:** Apply in a minimum finished spray volume of 100 L/ha by ground. Apply in a minimum finished spray volume of 50 L/ha by air.

SITE	PEST	RATE DUPONT™ CORAGEN™ insecticide /HA	SPECIFIC DIRECTIONS
<p>LEGUME VEGETABLES</p> <p>Bean (Lupinus) (includes grain lupin, sweet lupin, white lupin, and white sweet lupin);</p> <p>Bean (Phaseolus) (includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean);</p> <p>Bean (Vigna) (includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, yardlong bean);</p> <p>Broad bean (fava);</p> <p>Chickpea (garbanzo);</p> <p>Guar;</p> <p>Jackbean;</p> <p>Lablab bean;</p> <p>Lentil;</p> <p>Pea (Pisum) (includes dwarf pea, edible-podded pea, English pea, field pea, garden pea, green pea, snowpea,</p>	Grasshoppers	125 mL to 250 mL	<p>Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control.</p> <p>For cutworm control, apply to foliage when rain is not expected in the next 24 hours. For optimal control, apply to smaller plants or when lower portions of the plant can receive adequate coverage.</p>
	Cabbage looper Cutworms	250 mL	
	Armyworm Fall armyworm Beet armyworm Corn earworm European corn borer Western bean cutworm	250 mL to 375 mL	

sugar snap pea);			
Pigeon pea;			
Sword bean			
Soybean;			

**RESTRICTIONS AND PRECAUTIONS: Legume Vegetables**

- Do not make more than 4 applications per season.
- Do not apply more than once every 3 days.
- Do not apply less than 1 day before harvest. Observe a 1-day PHI.
- Do not exceed a total of 1.125 litres of DUPONT™ CORAGEN™ insecticide per ha per season.
- **Do not make a foliar application of DUPONT™ CORAGEN™ insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.**

**FRUITING VEGETABLES (EXCEPT CUCURBITS) – Crop Group 8**

**Spray Volume for Fruiting Vegetables:** Apply in a minimum finished spray volume of 100 L/ha by ground.

SITE	PEST	RATE DUPONT™ CORAGEN™ insecticide / ha	SPECIFIC DIRECTIONS
FRUITING VEGETABLES Eggplant, Groundcherry, Pepino, Pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper) Tomatillo, Tomato	Colorado potato beetle  Armyworm  Fall armyworm  Beet armyworm  Variegated cutworm  Tobacco hornworm  Tomato hornworm  Tomato fruitworm (Corn earworm)\  European corn borer	250 ml to 375 ml	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control.  For black cutworm control, apply to foliage when rain is not expected in the next 24 hours. For optimal control, apply to smaller plants or when lower portions of the plant can receive adequate coverage.  Colorado potato beetle resistance management: DO NOT apply DuPont™

	Cabbage looper Black cutworm	250 ml	CORAGEN™ insecticide for Colorado potato beetle control if any Group 28 Insecticide was used at planting as an in-furrow, soil or seed treatment.
--	---------------------------------	--------	---

**RESTRICTIONS AND PRECAUTIONS: Fruiting Vegetables**

- For pests with a range of application rates, use the high rate under heavy pest pressure.
- Do not make more than 4 applications per season.
- Do not apply more than once every 5 days.
- Do not apply less than 1 day before harvest. Observe a 1-day PHI.
- Do not exceed a total of 1.125 litres DUPONT™ CORAGEN™ insecticide per ha per season.
- **Do not make a foliar application of DUPONT™ CORAGEN™ insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.**

**CUCURBIT VEGETABLES – Crop Group 9**

**Spray Volume for Cucurbit Vegetables:** Apply in a minimum finished spray volume of 100 L/ha by ground.

SITE	PEST	RATE DUPONT™ CORAGEN™ insecticide / ha	SPECIFIC DIRECTIONS
CUCURBIT VEGETABLES Chayote (fruit); Chinese waxgourd (Chinese preserving	Cabbage looper Black cutworm	250 ml	Begin applications when treatment thresholds have been reached. Thorough coverage

melon); Citron melon; Cucumber; Gherkin; Gourd, edible (includes hyotan, cucuzza, hechima, Chinese okra); Momordica spp (includes balsam apple, balsam pear, bittermelon, Chinese cucumber); Muskmelon (includes cantaloupe); Pumpkin; Squash, summer; Squash, winter (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash); Watermelon	Armyworm  Fall armyworm  Corn earworm  Leafminers: <i>Liriomyza sativae</i> <i>Liriomyza trifolii</i>	250 ml to 375 ml	is important to obtain optimum control.  For black cutworm control, apply to foliage when rain is not expected in the next 24 hours. For optimal control, apply to smaller plants or when lower portions of the plant can receive adequate coverage.
---	---	------------------	--

#### RESTRICTIONS AND PRECAUTIONS: Cucurbit Vegetables

- For pests with a range of application rates, use the high rate under heavy pest pressure.
- Do not make more than 4 applications per season.
- Do not apply more than once every 5 days.
- Do not apply less than 1 day before harvest. Observe a 1-day PHI.
- Do not exceed a total of 1.125 litres DUPONT™ CORAGEN™ insecticide per ha per season.
- **Do not make a foliar application of DUPONT™ CORAGEN™ insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.**

#### CORN – FIELD, POP, SEED AND SWEET

**Spray Volume for Corn:** Apply in a minimum finished spray volume of 100 L/ha by ground. Apply in a minimum finished spray volume of 50 L/ha by air

SITE	PEST	RATE DUPONT™ CORAGEN™ insecticide / ha	SPECIFIC DIRECTIONS
CORN  Field corn; Pop corn; Seed corn.	Black cutworm	250 ml  Apply either by ground or aerial application equipment	Begin applications when treatment thresholds have been reached. Thorough coverage is important

Sweet Corn	Armyworm	250 ml to 375 ml	to obtain optimum control.
	Fall armyworm	Apply either by ground or aerial application equipment	For corn earworm, European corn borer or western bean cutworm control, time the applications to coincide with peak egg hatch. Scout by monitoring egg laying and egg hatch to determine application timing. Reapply if monitoring indicates it is necessary.
	Beet armyworm		
	Variegated cutworm		
	Corn earworm/ Tomato fruitworm		
	European corn borer		
	Western bean cutworm		
	Refer to the section of the label titled AERIAL APPLICATION LABEL INSTRUCTIONS for use information specific to aerial application.		

**RESTRICTIONS AND PRECAUTIONS: Corn**

- For pests with a range of application rates, use the high rate under heavy pest pressure.
- Do not make more than 4 applications per season.
- Do not apply more than once every 3 days on seed corn or sweet corn.
- Do not apply more than once every 7 days on field corn or popcorn.
- Do not apply less than 1 day before harvest of seed corn or sweet corn.
- Do not apply less than 14 days before harvest of field corn or popcorn.
- Do not exceed a total of 1.125 litres DUPONT™ CORAGEN™ insecticide per ha per season.
- **Do not make a foliar application of DUPONT™ CORAGEN™ insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.**

**CEREAL GRAINS – Crop Group 15 (EXCEPT CORN AND WILD RICE)**

**and CEREALS for FORAGE – Crop Group 16 (EXCEPT CORN AND WILD RICE)**

**Spray Volume for Cereal Grains:** Apply in a minimum finished spray volume of 100 L/ha by ground. Apply in a minimum finished spray volume of 50 L/ha by air

SITE	PEST	RATE DUPONT™ CORAGEN™ insecticide / ha	SPECIFIC DIRECTIONS
CEREAL GRAINS  Barley, Buckwheat, Millet, pearl Millet, proso, Oats, Rye, Sorghum, Teosinte, Triticale Wheat	Grasshoppers	125 mL to 250 mL	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control.
	Cutworms	250 mL	
	Armyworm	250 mL to 375 mL	
	Fall armyworm		
	Beet armyworm		
Corn earworm			
European corn borer			

**RESTRICTIONS AND PRECAUTIONS: Cereal Grains**

- Forage, Fodder and hay may be fed to livestock
- Do not make more than 3 applications per season.
- Do not apply more than once every 7 days.
- Do not apply less than 1 day before harvest. Observe a 1-day PHI
- Do not exceed a total of 1.125 litres DUPONT™ CORAGEN™ insecticide per ha per season
- For pests with a range of application rates, use the high rate under heavy pest pressure.
- **Do not make a foliar application of DUPONT™ CORAGEN™ insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.**

**GRASS FORAGE, FODDER, AND HAY GROUP – Crop Group 17**

**Spray Volume for Grass Forage, Fodder, and Hay Group:** Apply in a minimum finished spray volume of 100 L/ha by ground.

SITE	PEST	RATE DUPONT™ CORAGEN™ insecticide / ha	SPECIFIC DIRECTIONS
GRASS FORAGE FODDER AND HAY GROUP (Any grass,	Grasshoppers	125 mL to 250 mL	Begin applications when treatment thresholds have been

Gramineae family (either green or cured) except sugarcane and those included in the cereal grains group, that will be fed to or grazed by livestock, all pasture and range grasses and grasses grown for hay or silage	Armyworm Fall armyworm Beet armyworm Corn earworm	250 mL to 375 mL	reached. Thorough coverage is important to obtain optimum control.
--	--	------------------	--

**RESTRICTIONS AND PRECAUTIONS: Grass Forage, Fodder, and Hay Group**

- For any of the pests listed above, use the high rate under heavy pest pressure.
- Do not make more than 4 applications per season.
- Do not apply more than once every 7 days.
- Observe a 0-day PHI.
- Do not exceed a total of 1.125 litres DUPONT™ CORAGEN™ insecticide per ha per season.
- **Do not make a foliar application of DUPONT™ CORAGEN™ insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.**

**NON-GRASS ANIMAL FEEDS GROUP – Crop Group 18**

**Spray Volume for Non-Grass Animal Feeds Group:** Apply in a minimum finished spray volume of 100 L/ha by ground.

SITE	PEST	RATE DUPONT™ CORAGEN™ insecticide / ha	SPECIFIC DIRECTIONS
NON-GRASS ANIMAL FEEDS GROUP Alfalfa; Bean, velvet; Clover (Trifolium, Melilotus); Kudzu; Lespedeza; Lupin; Sainfoin; Trefoil; Vetch; Vetch, crown; Vetch, milk	Beet armyworm Alfalfa weevil (suppression only)	250 ml to 375 ml 375 ml to 500 ml	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control.

**RESTRICTIONS AND PRECAUTIONS: Non-Grass Animal Feeds Group**

- For either of the pests listed above, use the high rate under heavy pest pressure.
- Do not make more than 1 application per cutting.
- Observe a 0-day PHI.
- **Do not make a foliar application of DUPONT™ CORAGEN™ insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.**

## OILSEEDS GROUP – Crop Group 20

**Spray volume for Oilseeds Group:** Apply a minimum finished spray volume of 100 L/ha by ground. Apply a minimum finished spray volume of 50 L/ha by air.

SITE	PEST	RATE DUPONT™ CORAGEN™ Insecticide /ha	SPECIFIC DIRECTIONS
OILSEEDS GROUP  Canola Rape Seed; Mustard seed; Flax; Linseed; Sunflower seed; Safflower; Borage; Calendula; Crambe; Cuphea; Echium; Euphorbia; Evening Primrose; Gold of Pleasure; Hare's ear mustard; Lesquerella; Lunaria; Meadowfoam; Milkweed; Niger seed; Oil radish; Poppy seed; Rose hip; Stokes aster; Sweet rocket; Tea oil plant;	Diamondback moth	125	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control.
	Grasshoppers	125 - 250	
	Bertha armyworm	125 to 375	
	Cabbage looper	250	For early season cutworm control, apply to foliage when rain is not expected in the next 24 hours. For optimal control, apply to smaller plants or when lower portions of the plant can receive adequate coverage.
	Imported cabbage worm		
Swede midge			
Cutworms			
	Sunflower head moth	250 to 375	
	Reduces damage caused by banded sunflower moth.		

### RESTRICTIONS AND PRECAUTIONS: Oilseeds

- Do not make more than 3 applications per season.
- Do not apply more than once every 5 days.
- Do not apply less than 1 day before harvest. Observe a 1-day PHI
- Do not exceed a total of 1.125 litres DUPONT™ CORAGEN™ insecticide per ha per season
- **Do not make a foliar application of DUPONT™ CORAGEN™ insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.**

### MINT

**Spray Volume for Mint:** Apply in a minimum finished spray volume of 100 L/ha by ground.



SITE	PEST	RATE DUPONT™ CORAGEN™ insecticide / ha	SPECIFIC DIRECTIONS
MINT	Cabbage looper	250 ml	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control.

**RESTRICTIONS AND PRECAUTIONS: Mint**

- Do not make more than 4 applications per season.
- Do not apply more than once every 14 days.
- Do not apply less than 3 days before harvest. Observe a 3-day PHI.
- Do not exceed a total of 1 litre DUPONT™ CORAGEN™ insecticide per ha per season.
- **Do not make a foliar application of DUPONT™ CORAGEN™ insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.**

**OKRA**

**Spray Volume for Okra:** Apply in a minimum finished spray volume of 100 L/ha by ground.

SITE	PEST	RATE DUPONT™ CORAGEN™ insecticide / ha	SPECIFIC DIRECTIONS
OKRA	Beet armyworm  Corn earworm	250 ml to 375 ml	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control.

**RESTRICTIONS AND PRECAUTIONS: Okra**

- For either of the pests listed above, use the high rate under heavy pest pressure.
- Do not make more than 4 applications per season.
- Do not apply more than once every 5 days.
- Do not apply less than 1 day before harvest. Observe a 1-day PHI.
- Do not exceed a total of 1.125 litres DUPONT™ CORAGEN™ insecticide per ha per season.
- **Do not make a foliar application of DUPONT™ CORAGEN™ insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.**

**HOPS**

**Spray Volume for Hops:** Apply in a minimum finished spray volume of 100 L/ha by ground.

SITE	PEST	RATE DUPONT™	SPECIFIC DIRECTIONS
------	------	-----------------	---------------------

		CORAGEN™ Insecticide/HA	
Hops	Cutworms	250 to 375 ml	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control.
	Armyworm		
	Cabbage looper	250 mL	

**RESTRICTIONS AND PRECAUTIONS: Hops**

- Do not make more than 4 applications per season.
- Do not apply more than once every 7 days.
- Do not apply less than 0 day before harvest. Observe a 0 day PHI.
- Do not exceed a total of 1.125 litres of DUPONT™ CORAGEN™ insecticide per ha per season.
- Do not apply by air.

**NOTE TO USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT  
FOR SPECIAL USE APPLICATIONS:**

The DIRECTIONS FOR USE for this product for the uses described below were developed by persons other than The E. I. du Pont Canada Company and accepted for registration by Health Canada under the User Requested Minor Use Label Expansion program. The E. I. du Pont Canada Company itself makes no representation or warranty with respect to product performance (efficacy) or crop tolerance (phytotoxicity) claims for this product when used on the crops listed below. Accordingly, the Buyer and User assume all risks related to performance and crop tolerance arising, and agree to hold The E. I. du Pont Canada Company harmless from any claims based on efficacy or phytotoxicity in connection with these uses.

**FOR USE ON BRASSICA VEGETABLES - Crop Group 5**

**Spray Volume for Brassica Vegetables:** Apply in a minimum finished spray volume of 100 L/ha by ground.

SITE	PEST	RATE DUPONT™ CORAGEN™ insecticide/ ha	SPECIFIC DIRECTIONS
BRASSICA VEGETABLES Broccoli, Broccoli (Chinese), Broccoli raab, Brussels Sprouts, Cabbage, Cabbage (Chinese, bok choy), Cabbage (Chinese, napa), Cabbage (Chinese mustard, choy), Cauliflower, Cavalo broccolo, Collards, Kale, Kohlrabi, Mizuna, Mustard Greens, Mustard Spinach, Rape Greens	Swede midge	250 ml  For optimum control, apply Hasten NT Spray Adjuvant* at an application rate of 0.25% v/v or MSO Contentrate with Leci- Tech* at an application rate of 0.5% v/v.	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control.  Monitor Swede midge activity every 2-4 days using Jackson traps baited with pheromone lures.  Follow local threshold. Apply DUPONT™ Coragen™ insecticide as soon as possible once the Swede midge threshold is reached. Applications on cabbage, Chinese cabbage (napa), cauliflower and Chinese broccoli (gai lon) must be made early in the growing season, before heads start to form.  Given that there are multiple generations of Swede midge, several applications of insecticide (to a maximum of 4 per season) will likely be required.

**RESTRICTIONS AND PRECAUTIONS: Brassica Vegetables**

- Do not make more than 4 applications per season.
- Do not apply more than once every 3 days. For swede midge, make additional applications when monitoring indicates they are necessary, according to local thresholds.
- Do not apply less than 3 days before harvest. Observe a 3-day PHI.
- Do not exceed a total of 1 litre of DUPONT™ CORAGEN™ insecticide per ha per season.
- Do not apply by air.
- **Do not make a foliar application of DUPONT™ CORAGEN™ insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.**

**FOR USE ON Globe Artichokes**

**Spray Volume for Globe Artichoke:** Apply in a minimum finished spray volume of 100 L/ha by ground.

SITE	PEST	RATE DUPONT™ CORAGEN™ Insecticide/HA	SPECIFIC DIRECTIONS
Globe Artichoke	Cutworms	250 to 375 ml	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control.
	Armyworm		
	Cabbage looper	250 mL	

**RESTRICTIONS AND PRECAUTIONS: Artichokes**

- Do not make more than 4 applications per season.
- Do not apply more than once every 7 days.
- Do not apply less than 3 day before harvest. Observe a 3-day PHI.
- Do not exceed a total of 1.125 litres of DUPONT™ CORAGEN™ insecticide per ha per season.
- Do not apply by air.
- **Do not make a foliar application of DUPONT™ CORAGEN™ insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.**

**FOR USE ON GREENHOUSE CUCUMBERS, GREENHOUSE PEPPERS, GREENHOUSE EGGPLANTS, AND GREENHOUSE TOMATOES**

**Spray Volume for Greenhouse Vegetables:** For greenhouse tomatoes, the maximum finished spray volume is 1400 L/ha. For greenhouse cucumbers, greenhouse eggplants, and greenhouse peppers, the maximum finished spray volume is 2000 L/ha.

SITE	PEST	RATE DUPONT™ CORAGEN™ insecticide/1000 L finished spray volume	SPECIFIC DIRECTIONS
GREENHOUSE CUCUMBERS;  GREENHOUSE PEPPERS;  GREENHOUSE EGGPLANT;  GREENHOUSE TOMATOES	Cabbage looper	125 ml	Begin applications when treatment thresholds have been reached. Thorough coverage is required to obtain optimum control. Repeat applications if monitoring indicates it is necessary

**RESTRICTIONS AND PRECAUTIONS: Greenhouse Vegetables**

- Do not make more than 3 applications per crop cycle.
- Do not apply more than once every 7 days.
- Do not apply less than 1 day before harvest. Observe a 1-day PHI.

- Do not exceed a total of 750 ml of DUPONT™ CORAGEN™ insecticide per ha per crop cycle.
- Do not allow effluent or runoff from greenhouses containing this product to enter lakes, streams, ponds or other waters.
- **Do not make a foliar application of DUPONT™ CORAGEN™ insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.**

**FOR USE ON GREENHOUSE PEPPERS, GREENHOUSE EGGPLANTS, AND GREENHOUSE TOMATOES**

**Spray Volume for Greenhouse Vegetables:** Apply in a maximum finished spray volume of 1250 L/ha. Please note that the use of this product for control of tomato leafminer is restricted to plant growth stages for which thorough coverage can be achieved by application volumes of 1250 L/ha or less.

SITE	PEST	RATE DUPONT™ CORAGEN™ insecticide/1000 L finished spray volume	SPECIFIC DIRECTIONS
GREENHOUSE PEPPERS;  GREENHOUSE EGGPLANT;  GREENHOUSE TOMATOES	Lepidoptera n leafminer	200 ml / 1000L	Apply at egg hatch. Reapply if monitoring indicates it is necessary. Thorough coverage is important to obtain optimum control.

**RESTRICTIONS AND PRECAUTIONS: Greenhouse Vegetables**

- Do not make more than 3 applications per crop cycle.
- Do not apply more than once every 7 days.
- Do not apply less than 1 day before harvest. Observe a 1-day PHI.
- Do not exceed a total of 750 ml of DUPONT™ CORAGEN™ insecticide per ha per crop cycle.
- **Do not make a foliar application of DUPONT™ CORAGEN™ insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.**

**FOR USE ON PEANUTS**

**Spray Volume for Peanuts:** Apply in a minimum finished spray volume of 100 L/ha by ground.

SITE	PEST	RATE DUPONT™ CORAGEN™ Insecticide/HA	SPECIFIC DIRECTIONS
Peanuts	Armyworms  Cutworms  Corn	250 to 375 ml	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control.

	earworm		
--	---------	--	--

**RESTRICTIONS AND PRECAUTIONS: Peanuts**

- Do not make more than 4 applications per season.
- Do not apply by air.
- Do not apply more than once every 7 days.
- Do not apply less than 1 day before harvest. Observe a 1-day PHI.
- Do not exceed a total of 1.125 litres of DUPONT™ CORAGEN™ insecticide per ha per season.
- **Do not make a foliar application of DUPONT™ CORAGEN™ insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.**

**FOR USE ON GREEN ONIONS SUB GROUP – Crop group 3-07B**

**Spray Volume for Green Onion Subgroup:** Apply in a minimum finished spray volume of 100 L/ha by ground.

SITE	PEST	RATE DUPONT™ CORAGEN™ Insecticide/HA	SPECIFIC DIRECTIONS
Green onion sub group  Leek, Onion, green Onion, Welsh, tops, Shallot, fresh leaves, Chive, fresh leaves, Chive, Chinese, fresh leaves, Elegans hosta, Fritillaria, Kurrat, Lady's leek, Onion, Beltsville bunching, Onion, fresh, Onion, macrostem, Onion, tree, tops, Wild leek,	Cutworms  Leek moth (suppression )	250 to 375 ml	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control.

**RESTRICTIONS AND PRECAUTIONS: Green onion sub group**

- Do not make more than 4 applications per season.
- Do not apply by air.
- Do not apply more than once every 5 days.
- Do not apply less than 1 day before harvest. Observe a 1-day PHI.
- Do not exceed a total of 1.125 litres of DUPONT™ CORAGEN™ insecticide per ha per season.
- **Do not make a foliar application of DUPONT™ CORAGEN™ insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.**

## FIELD SPRAYER APPLICATION

**DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) fine classification. Boom height must be 60 cm or less above the crop or ground.

### Buffer zones:

Use of the following spray methods or equipment **DO NOT** require a buffer zone: hand-held or backpack sprayer and spot treatment.

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of Application	Crop	Buffer zones (meters) required for the protection of			
		Freshwater habitat of depths:		Estuarine/Marine Habitat of depths:	
		Less than 1m	Greater than 1 m	Less than 1 m	Greater than 1 m
Field sprayer	Root and Tuber vegetables, Fruiting vegetables, leafy vegetables, brassica vegetables, Legume vegetables cucurbit vegetables, corn, grass forage, fodder, and hay, non-grass animal feeds, mint, Oilseeds, okra, globe artichokes, peanuts, green onion subgroup and cereal grains	1	1	1	0

## AERIAL APPLICATION LABEL INSTRUCTIONS

**DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply when wind speed is greater than 16 km/h at flying height at the site of application. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) fine classification. To reduce drift caused by turbulent wingtip vortices, the nozzle distribution along the spray boom length **MUST NOT** exceed 65% of the wing- or rotor span.

Method of Application	Crop		Buffer zones (meters) required for the protection of			
			Freshwater habitat of depths:		Estuarine/Marine Habitat of depths:	
			Less than 1m	Greater than 1 m	Less than 1 m	Greater than 1 m
Aerial	Potato, Corn (field, pop, seed and sweet) Legume vegetables Oilseeds and cereal grains	Fixed Wing	15	1	1	0
		Rotary wing	10	1	1	0

Apply only by fixed-wing or rotary aircraft equipment, which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

#### Use Precautions for Aerial Application

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the “**National Aerial Pesticide Application Manual, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides**”.

#### Operator Precautions for Aerial Application

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking.

Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

#### Product Specific Precautions for Aerial Application

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-800-667-3925 or obtain technical advice from the distributor or your provincial agricultural representative. Application of this specific product must meet and/or conform to the following:

Apply the recommended rate in a minimum spray volume of 50 litres per hectare.

#### **MIXING INSTRUCTIONS:**

Spray equipment must be clean and free of previous pesticide deposits before applying DUPONT™ CORAGEN™ insecticide. Fill spray tank 1/4 to 1/2 full of water. Add DUPONT™ CORAGEN™ insecticide directly to spray tank. Mix thoroughly to fully disperse the insecticide; once dispersed



continued agitation is required. Once the product has been well mixed, add an adjuvant, where applicable, while continuously mixing. Recommendations regarding the use of adjuvants are provided under the Directions for Use Section of this label. Use mechanical or hydraulic means; do not use air agitation. Spray mix should not be stored overnight in spray tank.

#### **SPRAYER CLEANUP:**

Prior to application, start with clean, well maintained application equipment. Immediately following application, thoroughly clean all spray equipment to reduce the risk of forming hardened deposits which might become difficult to remove.

Drain spray equipment. Thoroughly rinse sprayer and flush hoses, boom and nozzles with clean water. Clean all other associated application equipment. Take all necessary safety precautions when cleaning equipment. Do not clean near wells, water sources or desirable vegetation. Dispose of waste rinse water in accordance with local regulations.

#### **PRECAUTIONS:**

- **KEEP OUT OF REACH OF CHILDREN.**
- Wear a long sleeved shirt, long pants and chemical resistant gloves while mixing /loading, applying and during clean up and repair.
- Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables are available, use detergent and hot water. Keep and wash PPE separately from other laundry.
- Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- For agricultural uses, do not enter or allow worker entry into treated areas during the restricted entry interval of 12 hours.
- See the Aerial Application Label Instructions section of this label for additional precautions specific to aerial application.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's website at [www.croplife.ca](http://www.croplife.ca).

#### **ENVIRONMENTAL HAZARDS:**

- Toxic to aquatic organisms. Observe buffer zones specified under DIRECTIONS FOR USE.
- Toxic to certain beneficial insects. Minimize spray drift to reduce harmful effects on beneficial insects in habitats next to the application site such as hedgerows and woodland.
- Chlorantraniliprole is persistent and may carryover. It is recommended that any products containing chlorantraniliprole not be used in areas treated with this product during the previous season.
- The use of this chemical may result in contamination of groundwater particularly in areas where soil is permeable (e.g. sandy soil) and/or the depth to the water table is shallow.
- To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.
- Avoid application when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.
- **DO NOT** apply this product directly to freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs, and wetlands), estuarine/marine habitats.
- **DO NOT** contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

#### **RESISTANCE-MANAGEMENT RECOMMENDATIONS:**

For resistance management, please note that DUPONT™ CORAGEN™ insecticide contains a Group 28 insecticide. Any insect population may contain individuals naturally resistant to

DUPONT™ CORAGEN™ insecticide and other Group 28 insecticides. The resistant individuals may dominate the insect population if this group of insecticides is used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but are specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance:

- Where possible, rotate the use of DUPONT™ CORAGEN™ insecticide or other Group 28 insecticides with different groups that control the same pests.
- Make no more than 2 successive applications per generation to the same insect species on a crop. The following application to the target pest(s) must be with an effective product with a different mode of action.
- Use tank mixtures with insecticides from a different group when such use is permitted.
- Insecticide use should be based on an IPM program that includes scouting and record keeping, and considers cultural, biological and other chemical control practices.
- Monitor treated pest populations for resistance development.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact DuPont Canada at 1-800-667-3925.

**STORAGE:**

Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed. To prevent contamination, store this product away from food or feed.

**DISPOSAL:**

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

**FIRST AID:**

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**IF SWALLOWED:** Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control

centre or doctor for further treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

For medical emergencies call 1-800-441-3637 (24 hours).

**TOXICOLOGICAL INFORMATION:** Treat symptomatically. This product contains a petroleum distillate. Vomiting may cause aspiration pneumonia.

**NOTICE TO USER:**

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

"®/TM Trademarks of E.I. du Pont de Nemours and Company, used under license. Member of CropLife Canada."

\*All other products mentioned are trademarks of their respective companies.