

PMRA Approved Label, A.P.
2015-2146
2015-10-27

GROUP	28	INSECTICIDE
--------------	-----------	--------------------

DUPONT™ Lumiderm™ Insecticide Seed Treatment

SUSPENSION

FOR SALE FOR USE ON CANOLA, RAPESEED, AND OILSEED MUSTARD
AGRICULTURAL

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

GUARANTEE: Cyantraniliprole 625 g/L

Contains 5-Chloro-2-methyl-2H-isothiazol-3-one at 0.00056% and 2-Methyl-2H-isothiazol-3-one at 0.00019%
as a preservative.

REGISTRATION NO 30894 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 100L, 1000L, BULK

For use in commercial seed treatment facilities only.

This product contains no colourant. An appropriate colourant must be added when this product is applied.
Regulations pertaining to the SEEDS ACT must be strictly adhered to when using this product. Seed must be
conspicuously coloured at the time of treatment.

E.I. DU PONT CANADA COMPANY
AGRICULTURAL PRODUCTS
P.O. BOX 2300, STREETSVILLE
MISSISSAUGA, ONTARIO
L5M 2J4
1-800-667-3925

PRECAUTIONS:

- **KEEP OUT OF REACH OF CHILDREN**
- Wear a long-sleeved shirt, long pants and chemical resistant gloves while mixing /loading, applying and during clean up and repair.
- Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables are available, use detergent and hot water. Keep and wash PPE separately from other laundry.
- Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Do not use treated seed for food, feed, or oil processing.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at: www.croplife.ca

FIRST AID:

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

For medical emergencies call 1-800-441-3637 (24 hours).

TOXICOLOGICAL INFORMATION: Treat symptomatically.

ENVIRONMENTAL HAZARDS:

- Toxic to aquatic organisms.
- Toxic to bees. This product is systemic. However bees are unlikely to be exposed to product residues in pollen and/or nectar resulting from seed treatment applications. When this product is applied and used according to label directions, risk to bees is expected to be negligible.
- **DO NOT** contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

LABELING OF TREATED SEED:

All bags containing treated seed must be labelled or tagged as follows:

- This seed has been treated with DuPont™ Lumiderm™ Insecticide Seed Treatment which contains cyantraniliprole.
- Do not use for feed, food or oil processing.
- Store away from feeds and other foodstuffs.
- Wear long-sleeved shirt, long pants and chemical-resistant gloves when handling treated seed.

- This product is toxic to aquatic organisms.
- Dispose of all excess treated seed. Left over treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements.
- Dispose of seed packaging in accordance with local requirements.
- Cover or incorporate spilled treated seeds.

STORAGE:

Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed. To prevent contamination store this product away from food or feed.

DISPOSAL:

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

FOR RECYCLABLE CONTAINERS:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use. If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

FOR RETURNABLE CONTAINERS:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

FOR REFILLABLE CONTAINERS:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

FOR DISPOSABLE CONTAINERS:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instruction for any required additional cleaning of the container prior to its disposal.
3. Make the empty, rinsed container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

<<DFU Booklet>>

GROUP	28	INSECTICIDE
--------------	-----------	--------------------

DUPONT™ Lumiderm™ Insecticide Seed Treatment

SUSPENSION

FOR SALE FOR USE ON CANOLA, RAPESEED, AND OILSEED MUSTARD

AGRICULTURAL

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

GUARANTEE: Cyantraniliprole 625 g/L

Contains 5-Chloro-2-methyl-2H-isothiazol-3-one at 0.00056% and 2-Methyl-2H-isothiazol-3-one at 0.00019% as a preservative.

REGISTRATION NO 30894 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 100L, 1000L, BULK

For use in commercial seed treatment facilities only.

This product contains no colourant. An appropriate colourant must be added when this product is applied. Regulations pertaining to the SEEDS ACT must be strictly adhered to when using this product. Seed must be conspicuously coloured at the time of treatment.

E.I. DU PONT CANADA COMPANY
AGRICULTURAL PRODUCTS
P.O. BOX 2300, STREETSVILLE
MISSISSAUGA, ONTARIO
L5M 2J4
1-800-667-3925

GENERAL INFORMATION

DuPont™ Lumiderm™ Insecticide Seed Treatment is a flowable suspension that is applied as a seed treatment for early season protection from flea beetle feeding damage for 28 to 35 days and early season protection from cutworm feeding damage.

DIRECTIONS FOR USE

This product contains no colourant. An appropriate colourant must be added when this product is applied. Regulations pertaining to the SEEDS ACT must be strictly adhered to when using this product. Seed must be conspicuously coloured at the time of treatment.

DuPont™ Lumiderm™ Insecticide Seed Treatment is an insecticide seed treatment for use in commercial seed treatment facilities only. It is not for use in on-farm treating systems such as hopper-box or slurry-box applications just prior to planting.

Apply only in areas with adequate ventilation or in areas that are equipped to remove mist or dust.

Use only a closed treatment system.

DO NOT use open cab planting equipment when planting canola seed treated with DuPont™ Lumiderm™ Insecticide Seed Treatment. Use only closed cab planting equipment. Workers working outside the closed cab must wear coveralls over a long-sleeved shirt, long pants, a dust mask, goggles, chemical-resistant gloves, socks and chemical-resistant boots.

CANOLA, RAPESEED, OILSEED MUSTARD

SITE	PEST	RATE mL of DuPont™ Lumiderm™ Insecticide Seed Treatment per 100 kg seed	SPECIFIC DIRECTIONS
CANOLA, RAPESEED, OILSEED MUSTARD	Flea beetles	960 to 1600	The application rates for flea beetles will also provide early season protection from cutworm feeding damage.
	Cutworms	480 to 960	Use the higher rates in areas with high pest pressure, or where extended early season control is required.

Do not make a subsequent foliar application of any Group 28 insecticide for a minimum of 60 days after planting seed treated with DuPont™ Lumiderm™ Insecticide Seed Treatment.

TANKMIX COMBINATIONS WITH COMMERCIAL SEED TREATMENTS

DUPONT™ LUMIDERM™ INSECTICIDE Seed Treatment may be tank mixed with Prosper^{EverGol} Seed Treatment, Helix Xtra Seed Treatment or Helix Vibrance Seed Treatment at labelled rates for early season control of flea beetles , early season protection from cutworm feeding damage, and control of diseases on the recommended tank mix partner labels.

SEED TREATMENT	RATE (mL product per 100 kg seed)	TARGET PESTS
DUPONT™ LUMIDERM™ INSECTICIDE Seed Treatment	640	Cutworms Flea beetles Seed and seedling diseases listed on tank mix partner labels
+ Prosper ^{EverGol} Seed Treatment	+ 1400	
OR	OR	
Helix Xtra Seed Treatment	1500	
OR	OR	
Helix Vibrance Seed Treatment	1500	

When tank mixing DUPONT™ LUMIDERM™ INSECTICIDE Seed Treatment, users must follow label instructions for both DUPONT™ LUMIDERM™ INSECTICIDE Seed Treatment and the tank mix partner.

In some cases, tank mixing a pest control product with another pest control product or a fertilizer can result in biological effects that could include, but are not limited to: reduced pest efficacy or increased host crop injury. The user assumes the risk of losses that result from the use of tank mixes that do not appear on this label or that are not specifically recommended by E. I. DuPont Canada Company.

Seed Storage:

Field and laboratory tests have demonstrated that application of DuPont™ Lumiderm™ Insecticide Seed Treatment to canola, rapeseed, and mustard seed will not negatively affect germination. However, due to seed quality and seed storage conditions beyond the control of E. I. du Pont Canada Company, no claims are made to guarantee the germination of carry-over seed.

Treatment of damaged seed, or seed known to be of low vigour and poor quality, may result in reduced germination and/or seed and seedling vigour. In cases where seed quality is unknown, treat a small portion of the seed with DuPont™ Lumiderm™ Insecticide Seed Treatment and confirm acceptable germination, prior to treating the entire seed lot.

MIXING INSTRUCTIONS:

Dilute in a sufficient volume to obtain thorough, uniform coverage.

Polymers, colourants, and other additives should be tested for compatibility and seed safety prior to use in combination with DuPont™ Lumiderm™ Insecticide Seed Treatment.

PRECAUTIONS:

- **KEEP OUT OF REACH OF CHILDREN.**
- Wear a long-sleeved shirt, long pants and chemical resistant gloves while mixing /loading, applying and during clean up and repair.
- Follow manufacturer’s instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables are available, use detergent and hot water. Keep and wash PPE separately from other laundry.
- Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Do not use treated seed for food, feed, or oil processing.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at: www.croplife.ca

FIRST AID:

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

For medical emergencies call 1-800-441-3637 (24 hours).

ENVIRONMENTAL HAZARDS:

- Toxic to aquatic organisms.
- Toxic to bees. This product is systemic. However bees are unlikely to be exposed to product residues in pollen and/or nectar resulting from seed treatment applications. When this product is applied and used according to label directions, risk to bees is expected to be negligible.
- **DO NOT** contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

ROTATIONAL CROP RESTRICTIONS:

Recommended Plant-Back Intervals (PBI) for Rotational Crops.

PBI	CROPS
0 days	Crop Group 1C (Tuberous and corm vegetables); Crop Group 3-07 (Bulb vegetables); Crop Group 4 (Leafy vegetables; except <i>Brassica</i> vegetables); Crop Group 5 (<i>Brassica</i> (Cole) leafy vegetables); Crop Group 8-09 (Fruiting vegetables); Crop Group 9 (Cucurbit vegetables); Crop Group 13-07B (Bushberry subgroup); Crop Group

	20 (Oilseed)
30 days	Crop Group 1A (Root and tuber vegetables); Crop Group 2 (Leaves of root and tuber vegetables); Crop Group 6 (Legume vegetables (succulent or dried); Crop Group 7 (Foliage of legume vegetables); Crop Group 13-07G (Lowgrowing berry subgroup); Crop Group 15 (Cereal grains); Crop Group 16 (Forage, fodder, and straw of cereal grains); Crop Group 17 (Grass forage, fodder, and hay); Crop Group 18 (Nongrass animal feeds: Forage, fodder, straw and hay)
365 days	Other crops

RESISTANCE-MANAGEMENT RECOMMENDATIONS:

For resistance management, please note that DuPont™ Lumiderm™ Insecticide Seed Treatment contains a Group 28 insecticide. Any insect population may contain individuals naturally resistant to DuPont™ Lumiderm™ Insecticide Seed Treatment and other Group 28 insecticides. The resistant individuals may dominate the insect population if this group of insecticides is used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but are specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance:

- Where possible, rotate the use of DuPont™ Lumiderm™ Insecticide Seed Treatment or other Group 28 insecticides with different groups that control the same pests.
- Use tank mixtures with insecticides from a different group when such use is permitted.
- Insecticide use should be based on an IPM program that includes scouting and record keeping, and considers cultural, biological and other chemical control practices.
- Monitor treated pest populations for resistance development.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact DuPont Canada at 1-800-667-3925.

STORAGE:

Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed. To prevent contamination store this product away from food or feed.

DISPOSAL:

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

FOR RECYCLABLE CONTAINERS:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use. If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

FOR RETURNABLE CONTAINERS:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

FOR REFILLABLE CONTAINERS:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

FOR DISPOSABLE CONTAINERS:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instruction for any required additional cleaning of the container prior to its disposal.
3. Make the empty, rinsed container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

"®/TM Trademarks of E.I. du Pont de Nemours and Company, used under license. Member of CropLife Canada."

*All other products mentioned are trademarks of their respective companies.